

Article

Group Technology Scheduling with Due-Date Assignment and Controllable Processing Times

Weiguo Liu * and Xuyin Wang

Business School, Northwest Normal University, Lanzhou 730070, China

* Correspondence: winterplum.home@nwnu.edu.cn or lwgwinterplum@163.com

Abstract: This paper investigates common (slack) due-date assignment single-machine scheduling with controllable processing times within a group technology environment. Under linear and convex resource allocation functions, the cost function minimizes scheduling (including the weighted sum of earliness, tardiness, and due-date assignment, where the weights are position-dependent) and resource-allocation costs. Given some optimal properties of the problem, if the size of jobs in each group is identical, the optimal group sequence can be obtained via an assignment problem. We then illustrate that the problem is polynomially solvable in $O(\varphi^3)$ time, where φ is the number of jobs.

Keywords: scheduling; group technology; position-dependent weights; single-machine; controllable processing times

1. Introduction

Classical scheduling problems consider fixed job processing times. However, scheduling problems with controllable processing times (\widetilde{CPT} , i.e., resource allocation) have received extensive attention (see Lu et al. [1], Liu et al. [2]). In 2018, Li and Wang [3] studied single-machine scheduling with deteriorating jobs and \widetilde{CPT} . For the general linear deterioration function, they proved that the weighted sum minimization of the makespan and total resource consumption costs can be solved in polynomial time. Lu and Liu [4] delved into single-machine scheduling with \widetilde{CPT} and position-dependent workloads. For scheduling and total resource consumption costs, they performed bicriterion analysis for the problem. In 2019, Geng et al. [5], and Sun et al. [6] investigated two-machine flow-shop problems with learning effects and \widetilde{CPT} . Under common due-date assignment and no-wait constraints, Geng et al. [5] proved that irregular objective minimization is solved in polynomial time. Under slack due-date assignment and no-wait constraints, Sun et al. [6] proved that irregular objective minimization is solved in polynomial time. In 2020, Liu and Jiang [7] studied scheduling with learning effects and \widetilde{CPT} on a two-machine no-wait flow-shop setting. Under common and slack due date assignments, they provided bicriterion analysis for scheduling and resource-consumption costs. In 2021, Lu et al. [8] considered a single-machine due-date assignment problem with \widetilde{CPT} and learning effects. Zhao [9], and Lv and Wang [10] revisited no-wait flow-shop problems with learning effects and \widetilde{CPT} . Under a slack (different) due-window assignment, Zhao [9] (Lv and Wang [10]) performed bicriterion analysis of scheduling (including earliness–tardiness penalties, due-window starting times, and the due-window size of all jobs) and resource-consumption costs. Zhao [9], and Lv and Wang [10] proved that several scheduling and resource-consumption costs can be solved in polynomial time. In 2022, Tian [11] addressed single-machine due-window assignment scheduling with \widetilde{CPT} . Under linear and convex resource allocation functions, the objective is to minimize generalized earliness and tardiness penalties. For common and slack due-window assignments, they demonstrated that the problem could be solved in polynomial time. In 2023, Wang et al. [12] explored single-machine scheduling with



Citation: Liu, W.; Wang, X. Group Technology Scheduling with Due-Date Assignment and Controllable Processing Times. *Processes* **2023**, *11*, 1271. <https://doi.org/10.3390/pr11041271>

Academic Editor: Luis Puigjaner

Received: 5 March 2023

Revised: 23 March 2023

Accepted: 18 April 2023

Published: 19 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

\widetilde{CPT} . Under linear and convex resource allocation functions, the objective was to minimize the weighted sum of general earliness–tardiness and resource-consumption costs where weights are position-dependent. They demonstrated that the problem was polynomially solvable.

In addition, the study of group technology (*GT*) is very important (see Liu [13]). In 2018, Wang et al. [14] considered single-machine scheduling with shortened job processing times. Under *GT* and ready times, they proved that some special cases of the makespan minimization could be solved in polynomial time. In 2019, Huang [15] scrutinized the scheduling with deteriorating jobs and *GT*, and proved that bicriterion single-machine minimization is polynomially solvable, where primary (secondary) criterion is the total weighted completion time (maximal cost). Liu et al. [16] focused on single-machine scheduling *GT* and deterioration effects. For makespan minimization with ready times, they proposed heuristic and branch-and-bound algorithms, and tested them via randomly instances. In 2021, Wang et al. [17] examined single-machine scheduling with *GT* and due-date assignment. For common, slack, and different due-date assignments, they proved that irregular objective minimization could be solved in $O(n \log n)$ time, where n is the number of jobs. In 2022, Wang et al. [18] investigated single-machine scheduling with *GT* and a shortened proportional linear processing time. For the general problem of makespan minimization, they proposed a heuristic algorithm and a branch-and-bound algorithm to solve the problem. In 2023, Chen et al. [19] scrutinized the single-machine problem with *GT* and a controllable learning effect. In due-window assignments, the objective is to minimize the total cost comprising due-window related penalties and investment costs. They proved that the problem could be solved in polynomial time.

To our knowledge, scheduling with *GT* and \widetilde{CPT} are concurrently widely reflected in real production (see Shabtay et al. [20], Zhu et al. [21], Wang et al. [22]). Wang and Liang [23], and Liang et al. [24] explored single-machine scheduling with *GT*, convex \widetilde{CPT} , and deterioration effects. In 2023, Yan et al. [25] studied single-machine scheduling with *GT* and \widetilde{CPT} . Under learning effects and limited resource availability, the goal is to minimize the total completion time. These authors proved that some special cases of the problem could be solved in polynomial time. For a general case of the problem, they also proposed heuristic and branch-and-bound algorithms. Chen et al. [26] worked on single-machine scheduling with *GT* and \widetilde{CPT} . In different due-date assignments and for a special case, they proved that the problem could be solved in polynomial time. In view of the importance of *GT* and \widetilde{CPT} , in this article, we continue the work of the concurrent single-machine scheduling with *GT* and \widetilde{CPT} for a common due-date assignment (Ξ_{CON} ; for details, see Gordon et al. [27]) and slack due-date assignment (Ξ_{SLK} ; see Gordon et al. [28], Liu et al. [29]). Our objective is to minimize the sum of scheduling (including the weighted sum of earliness, tardiness, and due-date assignments where weights are position-dependent (\widetilde{PDW} ; see Wang et al. [30], and Wang et al. [31])) and resource-allocation costs. This paper's contributions are as follows:

- We scrutinize the single-machine due-date assignment problem with the group technology and controllable processing times.
- Under Ξ_{CON} , and Ξ_{SLK} , the goal is to minimize the sum of scheduling (including the weighted sum of earliness, tardiness, and due-date assignment, where weights are \widetilde{PDW}) and resource-allocation costs.
- The optimal properties of a special case are presented, and we prove that the problem could be solved in polynomial time.

The rest of this article is organized as follows: In Section 2, we present the model. In Sections 3 and 4, we analyze the linear and convex resource functions, respectively. In Section 5, a numerical example is presented. In Section 6, we conclude the paper.

2. Problem Formulation

A set of φ jobs to be processed on a single-machine are divided into \aleph groups $\widetilde{G}_1, \widetilde{G}_2, \dots, \widetilde{G}_\aleph$, where a number of jobs belong to group \widetilde{G}_i is φ_i and $\varphi_1 + \varphi_2 + \dots + \varphi_\aleph = \varphi$. All jobs and the machine are available at Time 0. Machine setup time \widehat{s}_i is incurred before the jobs are processed in \widetilde{G}_i . There is not setup time between jobs in the same group, and jobs within each group must be processed consecutively. Let J_{ih} be the h th job in \widetilde{G}_i , $i = 1, \dots, \aleph; h = 1, \dots, \varphi_i$. For a linear resource function, the actual processing time of J_{ih} is

$$p_{ih} = a_{ih} - b_{ih}u_{ih}, 0 \leq u_{ih} \leq \bar{u}_{ih} < \frac{a_{ih}}{b_{ih}}, \tag{1}$$

where a_{ih} and b_{ih} are the normal processing time and positive compression rate of job J_{ih} , respectively (the normal processing time means that the processing time without any resource allocation), u_{ih} is the amount of a nonrenewable resource allocated to J_{ih} , and \bar{u}_{ih} denotes the maximal amount of the resource allocated to J_{ih} . For a convex resource function,

$$p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \tag{2}$$

where θ_{ih} is a workload of J_{ih} ($\ell > 0$ is a given constant).

Let C_{ih} and d_{ih} be the completion time and due date, respectively, of J_{ih} in \widetilde{G}_i . For the Ξ_{CON} assignment, we assumed that $d_{ih} = d_i$; for the Ξ_{SLK} assignment, we assumed that $d_{ih} = p_{ih} + q_i$, where q_i denotes common flow allowance for \widetilde{G}_i . Let $E_{ih} = \max\{0, d_{ih} - C_{ih}\}$ and $T_{ih} = \max\{0, C_{ih} - d_{ih}\}$ denote the earliness and tardiness, respectively, of job J_{ih} . Let $[r]$ be a scheduled job (group) in the r th position, $J_{i[r]}$ a scheduled job in the r th position in \widetilde{G}_i . Our goal was to find group sequence q and internal job sequence φ_i within \widetilde{G}_i , the set of due-dates $\vec{d} = \{d_1, d_2, \dots, d_m\}$ (flow allowances $\vec{q} = \{q_1, q_2, \dots, q_m\}$) and the resource allocation $\vec{u} = \{u_{ih} | i = 1, \dots, \aleph; h = 1, 2, \dots, \varphi_i\}$ such that cost function

$$\begin{aligned} H(q, \varphi_i | i = 1, \dots, \aleph, \vec{d} / \vec{q}, \vec{u}) &= \sum_{i=1}^{\aleph} \sum_{h=1}^{\varphi_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + \gamma d_i / q_i) + \sum_{i=1}^{\aleph} \sum_{h=1}^{\varphi_i} g_{ih}u_{ih} \\ &= \sum_{i=1}^m \sum_{h=1}^{\varphi_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i) \end{aligned} \tag{3}$$

is minimized, where α_{ih} and β_{ih} are position-dependent weights for earliness and tardiness costs, i.e., α_{ih} and β_{ih} are not related to job J_{ih} , but to position h in group \widetilde{G}_i , $\gamma \geq 0$ is a given constant, and g_{ih} is the cost of one unit of the allocated resource to job J_{ih} . With three field notations, this problem is denoted as follows:

$$1 \left| p_{ih} = a_{ih} - b_{ih}u_{ih}, \Xi_{CON} / \Xi_{SLK}, GT, \widehat{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\varphi_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i) \tag{4}$$

and

$$1 \left| p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \Xi_{CON} / \Xi_{SLK}, GT, \widehat{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\varphi_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i) \tag{5}$$

where 1 stands for the single-machine, field $\{p_{ih}, \Xi_{CON} / \Xi_{SLK}, GT, \widehat{PDW}\}$ denotes the characteristics of jobs and groups, and $\sum_{i=1}^{\aleph} \sum_{h=1}^{\varphi_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i)$ is the cost function. The notations and symbols used in this article are listed in Table 1.

Table 1. Symbols.

Symbol	Definition
\wp (resp. \aleph)	number of jobs (resp. groups)
\tilde{G}_i	group $i, i = 1, \dots, \aleph$
\wp_i	Number of jobs in \tilde{G}_i
J_{ih}	Job h in \tilde{G}_i
a_{ih} and b_{ih}	Normal processing time and compression rate, respectively, of J_{ih}
θ_{ih}	Workload of J_{ih}
u_{ih} and \bar{u}_{ih}	Amount and maximal amount, respectively, of the assigned resource to J_{ih}
p_{ih}	Actual processing time of J_{ih}
\tilde{s}_i	setup time of \tilde{G}_i
C_{ih} and d_{ih}	Completion time and due date, respectively, of J_{ih}
E_{ih} and T_{ih}	Earliness and tardiness, respectively, of J_{ih}
d_i and q_i	Common due date and flow allowance, respectively, of \tilde{G}_i
α_{ih} and β_{ih}	Position-dependent weights of earliness and tardiness, respectively, in the h th position in \tilde{G}_i
g_{ih}	Unit resource cost for J_{ih}

3. Linear Resource Function

There exists an optimal sequence that does not include idle machine times. Let \tilde{S}_i be the starting time of \tilde{G}_i , for a given job sequence φ_i within \tilde{G}_i ; completion times of $J_{i[h]}$ is

$$C_{i[h]} = \tilde{S}_i + \tilde{s}_i + \sum_{l=1}^h p_{i[l]} \quad (6)$$

Lemma 1. For a given job sequence φ_i and resource allocation within group \tilde{G}_i , under Ξ_{CON} and Ξ_{SLK} assignments, if the values of d_i and q_i , respectively, are within the starting and ending times of \tilde{G}_i , there exists an optimal value at which d_i and q_i are equal to the completion time of some job ($i = 1, \dots, \aleph$).

Proof. For the Ξ_{CON} assignment, it was assumed that $C_{i[k_i]} < d_i < C_{i[k_i+1]}$, where k_i is the k_i th position of group \tilde{G}_i , we have

$$\begin{aligned} H_i &= \sum_{h=1}^{\wp_i} (\alpha_{ih} E_{i[h]} + \beta_{ih} T_{i[h]} + g_{ih} u_{ih} + \gamma d_i) \\ &= \sum_{h=1}^{k_i} \alpha_{ih} E_{i[h]} + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} T_{i[h]} + \sum_{h=1}^{\wp_i} g_{ih} u_{ih} + \gamma \wp_i d_i \\ &= \sum_{h=1}^{k_i} \alpha_{ih} (d_i - C_{i[h]}) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} (C_{i[h]} - d_i) + \sum_{h=1}^{\wp_i} g_{ih} u_{ih} + \gamma \wp_i d_i \end{aligned} \quad (7)$$

If $d_i = C_{i[k_i]}$,

$$H'_i = \sum_{h=1}^{k_i} \alpha_{ih} (C_{i[k_i]} - C_{i[h]}) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} (C_{i[h]} - C_{i[k_i]}) + \sum_{h=1}^{\wp_i} g_{ih} u_{ih} + \gamma \wp_i C_{i[k_i]} \quad (8)$$

If $d_i = C_{i[k_i+1]}$,

$$H''_i = \sum_{h=1}^{k_i} \alpha_{ih} (C_{i[k_i+1]} - C_{i[h]}) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} (C_{i[h]} - C_{i[k_i+1]}) + \sum_{h=1}^{\wp_i} g_{ih} u_{ih} + \gamma \wp_i C_{i[k_i+1]} \quad (9)$$

Then

$$\begin{aligned}
 H_i - H'_i &= \sum_{h=1}^{k_i} \alpha_{ih}(d_i - C_{i[k_i]}) + \sum_{h=k_i+1}^{\varphi_i} \beta_{ih}(C_{i[k_i]} - d_i) + \gamma\varphi_i(d_i - C_{i[k_i]}) \\
 &= (d_i - C_{i[k_i]}) \left(\gamma\varphi_i + \sum_{h=1}^{k_i} \alpha_{ih} - \sum_{h=k_i+1}^{\varphi_i} \beta_{ih} \right)
 \end{aligned}
 \tag{10}$$

and

$$\begin{aligned}
 H_i - H''_i &= \sum_{h=1}^{k_i} \alpha_{ih}(d_i - C_{i[k_i+1]}) + \sum_{h=k_i+1}^{\varphi_i} \beta_{ih}(C_{i[k_i+1]} - d_i) + \gamma\varphi_i(d_i - C_{i[k_i+1]}) \\
 &= (d_i - C_{i[k_i+1]}) \left(\gamma\varphi_i + \sum_{h=1}^{k_i} \alpha_{ih} - \sum_{h=k_i+1}^{\varphi_i} \beta_{ih} \right)
 \end{aligned}
 \tag{11}$$

Thus, if $C_{i[k_i]} < d_i < C_{i[k_i+1]}$ and $\gamma\varphi_i + \sum_{h=1}^{k_i} \alpha_{ih} - \sum_{h=k_i+1}^{\varphi_i} \beta_{ih} \geq 0$, we have $H_i - H'_i \geq 0$; if $\gamma\varphi_i + \sum_{h=1}^{k_i} \alpha_{ih} - \sum_{h=k_i+1}^{\varphi_i} \beta_{ih} \leq 0$, we have $H_i - H'_i \leq 0$, hence, we can see that d_i coincides with some job completion time of \tilde{G}_i .

For the Ξ_{SLK} method, this result can be similarly obtained. \square

Lemma 2. For a given job sequence φ_i within \tilde{G}_i , under Ξ_{CON}/Ξ_{SLK} assignment, there exists an optimal $d_i = C_{i[k_i]}/(q_i = C_{i[k_i-1]})$ where k_i satisfies the following inequality: $\sum_{h=k_i+1}^{\varphi_i} \beta_{ih} - \sum_{h=1}^{k_i} \alpha_{ih} \leq \gamma\varphi_i \leq \sum_{h=k_i}^{\varphi_i} \beta_{ih} - \sum_{h=1}^{k_i-1} \alpha_{ih}$.

Proof. For the Ξ_{CON} assignment, from Lemma 1, it was assumed that $d_i = C_{i[k_i]}$; we then have

$$H_i = \sum_{h=1}^{k_i} \alpha_{ih}(C_{i[k_i]} - C_{i[h]}) + \sum_{h=k_i+1}^{\varphi_i} \beta_{ih}(C_{i[h]} - C_{i[k_i]}) + \sum_{h=1}^{n_i} g_{ih}u_{ih} + \gamma\varphi_i C_{i[k_i]}
 \tag{12}$$

With the technique of small perturbations, if $d_i = C_{i[k_i]} + \varepsilon$ ($\varepsilon \gg 0$),

$$H'_i = \sum_{h=1}^{k_i} \alpha_{ih}(C_{i[k_i]} + \varepsilon - C_{i[h]}) + \sum_{h=k_i+1}^{\varphi_i} \beta_{ih}(C_{i[h]} - C_{i[k_i]} - \varepsilon) + \sum_{h=1}^{n_i} g_{ih}u_{ih} + \gamma\varphi_i(C_{i[k_i]} + \varepsilon)
 \tag{13}$$

if $d_i = C_{i[k_i]} - \varepsilon$,

$$H''_i = \sum_{h=1}^{k_i-1} \alpha_{ih}(C_{i[k_i]} - \varepsilon - C_{i[h]}) + \sum_{h=k_i}^{\varphi_i} \beta_{ih}(C_{i[h]} - C_{i[k_i]} + \varepsilon) + \sum_{h=1}^{n_i} g_{ih}u_{ih} + \gamma\varphi_i(C_{i[k_i]} - \varepsilon)
 \tag{14}$$

$$H_i - H'_i = -\varepsilon \left(\gamma\varphi_i + \sum_{h=1}^{k_i} \alpha_{ih} - \sum_{h=k_i+1}^{\varphi_i} \beta_{ih} \right) \leq 0
 \tag{15}$$

$$H_i - H''_i = \varepsilon \left(\gamma\varphi_i + \sum_{h=1}^{k_i-1} \alpha_{ih} - \sum_{h=k_i}^{\varphi_i} \beta_{ih} \right) \leq 0
 \tag{16}$$

Hence, k_i satisfies $\gamma\varphi_i + \sum_{h=1}^{k_i} \alpha_{ih} - \sum_{h=k_i+1}^{\varphi_i} \beta_{ih} \geq 0$ and $\gamma\varphi_i + \sum_{h=1}^{k_i-1} \alpha_{ih} - \sum_{h=k_i}^{\varphi_i} \beta_{ih} \leq 0$, i.e., $\sum_{h=k_i+1}^{\varphi_i} \beta_{ih} - \sum_{h=1}^{k_i} \alpha_{ih} \leq \gamma\varphi_i \leq \sum_{h=k_i}^{\varphi_i} \beta_{ih} - \sum_{h=1}^{k_i-1} \alpha_{ih}$.

For the Ξ_{SLK} assignment, the result can similarly be obtained. \square

Remark 1. If k_i does not satisfy inequality $\sum_{h=k_i+1}^{\varphi_i} \beta_{ih} - \sum_{h=1}^{k_i} \alpha_{ih} \leq \gamma\varphi_i \leq \sum_{h=k_i}^{\varphi_i} \beta_{ih} - \sum_{h=1}^{k_i-1} \alpha_{ih}$, we can set $k_i = 0$.

For the Ξ_{CON} assignment, from Equation (6), Lemma 2, and $d_i = C_{i[k_i]} = \tilde{S}_i + \tilde{s}_i + \sum_{l=1}^{k_i} p_{i[l]}$, we have

$$\begin{aligned}
 H(\Xi_{CON}) &= \sum_{i=1}^{\aleph} \left(\sum_{h=1}^{k_i} \alpha_{ih} (d_i - C_{i[h]}) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} (C_{i[h]} - d_i) + \gamma \wp_i d_i + \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \right) \\
 &= \sum_{i=1}^{\aleph} \left(\sum_{h=1}^{k_i} \alpha_{ih} (C_{i[k_i]} - C_{i[h]}) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} (C_{i[h]} - C_{i[k_i]}) + \gamma \wp_i C_{i[k_i]} + \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \right) \\
 &= \sum_{i=1}^{\aleph} \left(\sum_{h=1}^{k_i} \alpha_{ih} \left(\sum_{l=1}^{k_i} p_{i[l]} - \sum_{l=1}^h p_{i[l]} \right) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} \left(\sum_{l=1}^h p_{i[l]} - \sum_{l=1}^{k_i} p_{i[l]} \right) \right) \\
 &\quad + \gamma \sum_{i=1}^{\aleph} \wp_i \left(\tilde{S}_i + \tilde{s}_i + \sum_{l=1}^{k_i} p_{i[l]} \right) + \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \\
 &= \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} \zeta_{ih} p_{i[h]} + \gamma \sum_{i=1}^{\aleph} \wp_i (\tilde{S}_i + \tilde{s}_i) + \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \tag{17}
 \end{aligned}$$

where

$$\zeta_{ih} = \begin{cases} \sum_{l=1}^{h-1} \alpha_{il} + \gamma n_i, & h = 1, 2, \dots, k_i, \\ \sum_{l=h}^{k_i} \beta_{il}, & h = k_i + 1, k_i + 2, \dots, \wp_i. \end{cases} \tag{18}$$

For the Ξ_{SLK} assignment, from Equation (6), Lemma 2, and $q_i = C_{i[k_i-1]} = \tilde{S}_i + \tilde{s}_i + \sum_{l=1}^{k_i-1} p_{i[l]}$, we have

$$\begin{aligned}
 H(\Xi_{SLK}) &= \sum_{i=1}^{\aleph} \left(\sum_{h=1}^{k_i} \alpha_{ih} (d_{i[h]} - C_{i[h]}) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} (C_{i[h]} - d_{i[h]}) + \gamma \wp_i q_i + \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \right) \\
 &= \sum_{i=1}^{\aleph} \sum_{h=1}^{k_i} \alpha_{ih} (p_{i[h]} + C_{i[k_i-1]} - C_{i[h]}) + \sum_{i=1}^{\aleph} \sum_{h=k_i+1}^{\wp_i} \beta_{ih} (C_{i[h]} - p_{i[h]} - C_{i[k_i-1]}) \\
 &\quad + \sum_{i=1}^{\aleph} \gamma \wp_i C_{i[k_i-1]} + \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \\
 &= \sum_{i=1}^{\aleph} \left(\sum_{h=1}^{k_i} \alpha_{ih} \left(\sum_{l=1}^{k_i-1} p_{i[l]} - \sum_{l=1}^{h-1} p_{i[l]} \right) + \sum_{h=k_i+1}^{\wp_i} \beta_{ih} \left(\sum_{l=1}^{h-1} p_{i[l]} - \sum_{l=1}^{k_i-1} p_{i[l]} \right) \right) \\
 &\quad + \gamma \sum_{i=1}^{\aleph} \wp_i \left(\tilde{S}_i + \tilde{s}_i + \sum_{l=1}^{k_i-1} p_{i[l]} \right) + \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \\
 &= \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} \zeta_{ih} p_{i[h]} + \gamma \sum_{i=1}^{\aleph} \wp_i (\tilde{S}_i + \tilde{s}_i) + \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} g_{i[h]} u_{i[h]} \tag{19}
 \end{aligned}$$

where

$$\zeta_{ih} = \begin{cases} \sum_{l=1}^h \alpha_{il} + \gamma n_i, & h = 1, 2, \dots, k_i - 1, \\ \sum_{l=h+1}^{k_i} \beta_{il}, & h = k_i, k_i + 1, \dots, \wp_i - 1, \\ 0, & h = \wp_i. \end{cases} \tag{20}$$

Since k_i values are independent of ϱ and \wp_i ($i = 1, \dots, \aleph$), from Lemmas 1 and 2, if $p_{ih} = a_{ih} - b_{ih} u_{ih}$, the cost objective can be expressed:

$$\begin{aligned}
 H(\Xi_{CON}/\Xi_{SLK}) &= \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} a_{[i]h} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^{\aleph} (\wp_{[r]}) \right) + \gamma \sum_{i=1}^{\aleph} \left(\wp_{[i]} \times \sum_{r=1}^i \tilde{s}_{[r]} \right) \\
 &\quad + \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} \left(g_{[i]h} - b_{[i]h} \right) \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^{\aleph} (\wp_{[r]}) \right) u_{[i]h} \tag{21}
 \end{aligned}$$

where for the Ξ_{CON} assignment,

$$\zeta_{[i]h} = \begin{cases} \sum_{l=1}^{h-1} \alpha_{[i]l} + \gamma \wp_{[i]}, & h = 1, 2, \dots, k_i, \\ \sum_{l=h}^{k_i} \beta_{[i]l}, & h = k_i + 1, k_i + 2, \dots, \wp_{[i]}. \end{cases} \tag{22}$$

For the Ξ_{SLK} assignment,

$$\zeta_{[i]h} = \begin{cases} \sum_{l=1}^h \alpha_{[i]l} + \gamma \wp_{[i]}, & h = 1, 2, \dots, k_i - 1, \\ \sum_{l=h+1}^{\wp_{[i]}} \beta_{[i]l}, & h = k_i, k_i + 1, \dots, \wp_{[i]} - 1, \\ 0, & h = \wp_{[i]}. \end{cases} \tag{23}$$

Lemma 3. In the given group sequence and job sequence within each group, optimal resource allocation $\tilde{\mathbf{u}}^*(\varrho, \wp_1, \dots, \wp_N)$ is

$$u_{[i][h]}^* = \begin{cases} \bar{u}_{[i][h]}, & \text{if } g_{[i][h]} - b_{[i][h]} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(\wp_{[r]} \right) \right) < 0, \\ u_{[i][h]} \in [0, \bar{u}_{[i][h]}], & \text{if } g_{[i][h]} - b_{[i][h]} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(\wp_{[r]} \right) \right) = 0, \\ 0, & \text{if } g_{[i][h]} - b_{[i][h]} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(\wp_{[r]} \right) \right) > 0. \end{cases} \tag{24}$$

Proof. Let the derivative of Equation (21) with respect to $u_{[i][j]}$ be equal to 0, and the result can be obtained. \square

For a given group order ϱ , from Equation (21), $\sum_{i=1}^N \sum_{h=1}^{\wp_{[i]}} a_{[i][h]} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(\wp_{[r]} \right) \right) + \sum_{i=1}^N \sum_{h=1}^{\wp_{[i]}} \left(g_{[i][h]} - b_{[i][h]} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(\wp_{[r]} \right) \right) \right) u_{[i][h]}$ is dependent only on the internal job sequence, while term $\gamma \sum_{i=1}^N \left(\wp_{[i]} \times \sum_{r=1}^i \bar{s}_{[r]} \right)$ is independent of the internal job sequence within each group. Now, we prove that the optimal sequence within each group can be obtained with the following lemma.

Lemma 4. Given group order ϱ , optimal job sequence $\varphi_i^*(\varrho)$ within $\widetilde{G}_i(\varrho)$ is obtained in $O(\wp_i^3)$ time.

Proof. For a given $\widetilde{G}_{[i]}$, let $x_{jh}^{[i]}$ be a binary variable, i.e., if $J_{[i]j}$ in $\widetilde{G}_{[i]}$ is assigned to h th position, $x_{jh}^{[i]} = 1$; otherwise, $x_{jh}^{[i]} = 0, j, h = 1, \dots, \wp_{[i]}$. Let

$$\vartheta_{jh}^{[i]} = \begin{cases} g_{[i]j} \bar{u}_{[i]j} + (a_{[i]j} - b_{[i]j} \bar{u}_{[i]j}) \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(n_{[r]} \right) \right), & \text{if } g_{[i]j} - b_{[i]j} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(\wp_{[r]} \right) \right) \leq 0; \\ a_{[i]j} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(n_{[r]} \right) \right), & \text{if } g_{[i]j} - b_{[i]j} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^N \left(\wp_{[r]} \right) \right) > 0. \end{cases} \tag{25}$$

As in Wang et al. [22], optimal job sequence $\varphi_i^*(\varrho)$ can be obtained with the assignment problem (\widetilde{AP}):

$$\text{Min } \sum_{j=1}^{\wp_{[i]}} \sum_{h=1}^{\wp_{[i]}} \vartheta_{jh}^{[i]} x_{jh}^{[i]} \tag{26}$$

s.t.

$$\sum_{j=1}^{\wp_{[i]}} x_{jh}^{[i]} = 1, \quad h = 1, \dots, \wp_{[i]}, \tag{27}$$

$$\sum_{h=1}^{\wp_{[i]}} x_{jh}^{[i]} = 1, \quad j = 1, \dots, \wp_{[i]}, \tag{28}$$

$$x_{jh}^{[i]} \in \{0, 1\}, \quad j, h = 1, \dots, \wp_i. \tag{29}$$

The above \widetilde{AP} is solvable in $O(n_{[i]}^3)$ time; hence, determining the total complexity of $\varphi_i(\varrho)$ ($i = 1, \dots, \aleph$) is bounded by $\sum_{i=1}^{\aleph} O(\wp_i^3) = O(\wp^3)$. \square

For

$$1 \left| p_{ih} = a_{ih} - b_{ih}u_{ih}, \Xi_{CON} / \Xi_{SLK}, GT, PDW \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i),$$

from Lemmas 1–4, the complexity of determining the optimal group sequence is still an open problem, so we discuss a special case, i.e., $\wp_i = \widehat{N}$, $i = 1, \dots, \aleph$.

Lemma 5. For

$$1 \left| p_{ih} = a_{ih} - b_{ih}u_{ih}, \Xi_{CON} / \Xi_{SLK}, GT, PDW \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i),$$

if $\wp_i = \widehat{N}$ ($i = 1, \dots, \aleph$), the optimal group sequence ϱ^* is obtained by \widetilde{AP} in $O(\wp^3)$ time.

Proof. From Equation (21), cost function (3) is determined by both the group and job sequences. Optimal job sequence φ_i^* can be obtained with Lemma 4, and the cost function with φ_i^* is just dependent on the i th group position in ϱ . In $n_i = \widehat{N}$ ($i = 1, \dots, \aleph$), term $\gamma \sum_{i=1}^{\aleph} (\wp_{[i]} \times \sum_{r=1}^i \widetilde{s}_{[r]}) = \gamma \widehat{N} \sum_{i=1}^{\aleph} [(\aleph - i + 1) \widetilde{s}_{[i]}]$. Let y_{ir} be a binary variable. If group \widetilde{G}_i is assigned to r th position, $y_{ir} = 1$; otherwise, $y_{ir} = 0$, $i, r = 1, \dots, \aleph$. Let

$$\chi_{ir} = \begin{cases} \sum_{j=1}^{\widehat{N}} (g_{i[j]} \bar{u}_{i[j]} + (a_{i[j]} - b_{i[j]} \bar{u}_{i[j]}) (\xi_{ij} + \gamma(\aleph - r) \widehat{N})) + \gamma \widehat{N} (\aleph - r + 1) \widetilde{s}_i, & \text{if } g_{i[j]} - b_{i[j]} (\xi_{ij} + \gamma(\aleph - r) \widehat{N}) \leq 0; \\ \sum_{j=1}^{\widehat{N}} (a_{i[j]} (\xi_{ij} + \gamma(\aleph - r) \widehat{N})) + \gamma \widehat{N} (\aleph - r + 1) \widetilde{s}_i, & \text{if } g_{i[j]} - b_{i[j]} (\xi_{i[j]} + \gamma(\aleph - r) \widehat{N}) > 0. \end{cases} \tag{30}$$

As in Shabtay et al. [20], optimal group sequence ϱ^* was obtained with the following \widetilde{AP} :

$$\text{Min } \sum_{i=1}^{\aleph} \sum_{r=1}^{\aleph} \chi_{ir} y_{ir} \tag{31}$$

s.t.

$$\sum_{i=1}^{\aleph} y_{ir} = 1, \quad r = 1, \dots, \aleph, \tag{32}$$

$$\sum_{r=1}^{\aleph} y_{ir} = 1, \quad i = 1, \dots, \aleph, \tag{33}$$

$$y_{ir} \in \{0, 1\}, \quad i, r = 1, \dots, \aleph. \tag{34}$$

The above \widetilde{AP} is solvable in $O(\aleph^3) \leq O(\wp^3)$ time. \square

Via Lemmas 1–5 and the above analysis, for $\wp_i = \widehat{N}$ ($i = 1, \dots, \aleph$), the following algorithm (i.e., Algorithm 1) is presented to solve

$$1 \left| p_{ih} = a_{ih} - b_{ih}u_{ih}, \Xi_{CON} / \Xi_{SLK}, GT, \widetilde{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i).$$

Theorem 1. If $\wp_i = \widehat{N}$ ($i = 1, \dots, \aleph$), Algorithm 1 solves

$$1 \left| p_{ih} = a_{ih} - b_{ih}u_{ih}, \Xi_{CON} / \Xi_{SLK}, GT, \widetilde{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i)$$

in $O(\wp^3)$ time.

Algorithm 1: Linear resource function

Step 1. Calculate k_i by Lemma 2.

Step 2. For each possible position of each group in ϱ , calculate $\vartheta_{jh}^{[i]}$ with Equation (25) for $j, h = 1, \dots, \widehat{N}$, where for the Ξ_{CON} assignment, $\zeta_{[i]h}$ is given by Equation (22), and for the Ξ_{SLK} assignment, $\tilde{\zeta}_{[i]h}$ is given by Equation (23).

Step 3. Solve \widehat{AP} (26)–(29) to find internal job sequence φ_{ir}^* within \widetilde{G}_i if this group is assigned to the r th position in ϱ .

Step 4. Calculate χ_{ir} with Equation (30) with φ_{ir}^* for $i, r = 1, \dots, \aleph$.

Step 5. Solve \widetilde{AP} (31)–(34) to find optimal sequences ϱ^* and φ_i^* .

Step 6. Compute optimal resource allocation $u_{ih}^*(\varrho^*, \varphi_1^*, \dots, \varphi_{\aleph}^*)$ with Equation (24).

Step 7. For the Ξ_{CON} and Ξ_{SLK} assignments, calculate $d_i^* = C_{i[k_i]}$ and $q_i^* = C_{i[k_i-1]}$, respectively, with Lemma 2.

Proof. With Lemmas 1–5, the correctness of Algorithm 1 can be confirmed. Steps 1, 6, and 7 need $O(\wp)$ time; Steps 2 and 3 need $O(\wp^3)$ time; Steps 4 and 5 need $O(\aleph^3) \leq O(\wp^3)$ time. Thus, the total computational time is $O(\wp^3)$. □

4. Convex Resource Function

Similar to Section 3, for problem

$$1 \left| p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \Xi_{CON} / \Xi_{SLK}, GT, \widetilde{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i),$$

we have

$$H(\Xi_{CON} / \Xi_{SLK}) = \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^{\aleph} (\wp_{[r]}) \right) \left(\frac{\theta_{[i][h]}}{u_{[i][h]}} \right)^\ell + \gamma \sum_{i=1}^{\aleph} \left(\wp_{[i]} \times \sum_{r=1}^i \widetilde{s}_{[r]} \right) + \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} g_{[i][h]} u_{[i][h]}, \tag{35}$$

where for the Ξ_{CON} assignment, $\zeta_{[i]h}$ is given by Equation (22), and for the Ξ_{SLK} assignment, $\tilde{\zeta}_{[i]h}$ is given by Equation (23), $i, h = 1, \dots, \wp_i$.

Lemma 6. Under the given group and job sequences

$$1 \left| p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \Xi_{CON} / \Xi_{SLK}, GT, \widetilde{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i / q_i),$$

the optimal resource allocation $\tilde{\mathbf{u}}^*(\varrho, \varphi_1, \dots, \varphi_{\aleph})$ is

$$u_{[i][h]}^* = \left(\frac{\ell \left(\zeta_{[i]h} + \gamma \sum_{r=i+1}^{\aleph} (\wp_{[r]}) \right)}{g_{[i][h]}} \right)^{\frac{1}{\ell+1}} \times \left(\theta_{[i][h]} \right)^{\frac{\ell}{\ell+1}}, \tag{36}$$

where for the Ξ_{CON} assignment, $\xi_{[i]h}$ is given by Equation (22), and for the Ξ_{SLK} assignment, $\xi_{[i]h}$ is given by Equation (23), $i, h = 1, \dots, \wp_i$.

Proof. From Equation (35), H is a convex function of $u_{[i][h]}$; hence, let

$$\frac{\partial H}{\partial u_{[i][h]}} = g_{[i][h]} - \ell \left(\xi_{[i]h} + \gamma \sum_{r=i+1}^{\aleph} (\wp_{[r]}) \right) \frac{(\theta_{[i][h]})^\ell}{(u_{[i][h]})^{\ell+1}} = 0,$$

and the result of Equation (36) can be obtained. \square

By substituting Equation (36) into Equation (35), it follows that

$$\begin{aligned} H(\varrho, \varphi_1, \dots, \varphi_\aleph, \mathbf{u}^*) &= \left(\ell^{\frac{-\ell}{\ell+1}} + \ell^{\frac{1}{\ell+1}} \right) \sum_{i=1}^{\aleph} \sum_{j=1}^{\wp_{[i]}} \left(\xi_{[i]h} + \gamma \sum_{r=i+1}^{\aleph} (\wp_{[r]}) \right)^{\frac{1}{\ell+1}} (g_{[i][h]} \theta_{[i][h]})^{\frac{\ell}{\ell+1}} \\ &\quad + \gamma \sum_{i=1}^{\aleph} \left(\wp_{[i]} \times \sum_{r=1}^i \widetilde{s}_{[r]} \right) \end{aligned} \tag{37}$$

For the Ξ_{CON} assignment, let

$$\xi_{ih} = \begin{cases} \sum_{l=1}^{h-1} \alpha_{il} + \gamma \wp_i, & h = 1, \dots, k_i, \\ \sum_{l=h}^{n_i} \beta_{il}, & h = k_i + 1, k_i + 2, \dots, \wp_i, \end{cases} \tag{38}$$

For the Ξ_{SLK} assignment, let

$$\xi_{ih} = \begin{cases} \sum_{l=1}^h \alpha_{il} + \gamma n_{[i]}, & h = 1, \dots, k_i - 1, \\ \sum_{l=h+1}^{n_i} \beta_{il}, & h = k_i, k_i + 1, \dots, \wp_i - 1, \\ 0, & h = \wp_i. \end{cases} \tag{39}$$

Lemma 7. Given group order ϱ , optimal job sequence φ_i^* ($i = 1, \dots, \aleph$) within \widetilde{G}_i can be obtained by matching the smallest and second smallest ξ_{ih} to the job with the largest and second largest $g_{ih}\theta_{ih}$, respectively, and so on.

Proof. From Equation (37), $\left(\ell^{\frac{-\ell}{\ell+1}} + \ell^{\frac{1}{\ell+1}} \right)$ is a given constant, $\gamma \sum_{r=i+1}^{\aleph} (\wp_{[r]})$ and $\gamma \sum_{i=1}^{\aleph} \left(\wp_{[i]} \times \sum_{r=1}^i \widetilde{s}_{[r]} \right)$ are independent of the internal job sequence within each group. According to Hardy et al. [32], the optimal job sequence for \widetilde{G}_i is obtained by matching the smallest and second smallest ξ_{ih} to the job with the largest and second largest $g_{ih}\theta_{ih}$, respectively, and so on. \square

For

$$1 \left| p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \Xi_{CON}/\Xi_{SLK}, GT, \widetilde{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih} E_{i[h]} + \beta_{ih} T_{i[h]} + g_{ih} u_{ih} + \gamma d_i / q_i),$$

the complexity of finding the optimal group sequence is still an open problem, but for the special case $\wp_i = \widehat{N}$, $i = 1, \dots, \aleph$, the optimal ϱ^* is obtained in $O(\varphi^3)$ time.

Lemma 8. For

$$1 \left| p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \Xi_{CON}/\Xi_{SLK}, GT, \widetilde{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih} E_{i[h]} + \beta_{ih} T_{i[h]} + g_{ih} u_{ih} + \gamma d_i / q_i),$$

if $\varphi_i = \widehat{N}$ ($i = 1, \dots, \aleph$), the optimal group sequence q^* can be determined with \widetilde{AP} in $O(\varphi^3)$ time.

Proof. Similar to Lemma 5, from Equation (37), let

$$\chi_{ir} = \left(\ell^{\frac{-\ell}{\ell+1}} + \ell^{\frac{1}{\ell+1}} \right) \sum_{h=1}^{\widehat{N}} \left(\xi_{ih} + \gamma \widehat{N}(\aleph - r) \right)^{\frac{1}{\ell+1}} \left(g_{i[h]} \theta_{i[h]} \right)^{\frac{\ell}{\ell+1}} + \gamma \widehat{N}(\aleph - r + 1) \widetilde{s}_i, \quad (40)$$

The optimal group sequence q^* can be obtained with \widetilde{AP} (31)–(34), where χ_{ir} is given by Equation (40). \square

Similarly, for $n_i = \widehat{N}$ ($i = 1, \dots, \aleph$), the following algorithm (i.e., Algorithm 2) is presented to solve

$$1 \mid p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \text{CON, GT, PDW} \left| \sum_{i=1}^{\aleph} \sum_{h=1}^{\varphi_i} (\alpha_{ih} E_{i[h]} + \beta_{ih} T_{i[h]} + g_{ih} u_{ih} + \gamma d_i).$$

Theorem 2. If $\varphi_i = \widehat{N}$ ($i = 1, \dots, \aleph$), Algorithm 2 solves

$$1 \mid p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \Xi_{\text{CON}} / \Xi_{\text{SLK}}, \text{GT, PDW} \left| \sum_{i=1}^{\aleph} \sum_{h=1}^{\varphi_i} (\alpha_{ih} E_{i[h]} + \beta_{ih} T_{i[h]} + g_{ih} u_{ih} + \gamma d_i / q_i)$$

in $O(\varphi^3)$ time.

Algorithm 2: Convex resource function

Step 1. Calculate k_i by Lemma 2.

Step 2. For each group \widetilde{G}_i ($i = 1, \dots, \aleph$), Lemma 7 is used to obtain internal job sequence φ_i^* , where for the Ξ_{CON} assignment, ξ_{ih} is given by Equation (38), and for the Ξ_{SLK} assignment, ξ_{ih} is given by Equation (39), $i = 1, \dots, \aleph, h = 1, \dots, \varphi_i$.

Step 3. χ_{ir} is computed with Equation (40) with φ_i^* for $i, r = 1, \dots, \aleph$.

Step 4. Solve \widetilde{AP} (31)–(34) to determine the optimal group sequence q^* .

Step 5. Compute optimal resource allocation $u_{ih}^*(q^*, \varphi_1^*, \dots, \varphi_\aleph^*)$ with Equation (36).

Step 6. For the Ξ_{CON} and Ξ_{SLK} assignments, calculate $d_i^* = C_{i[k_i]}$ and $q_i^* = C_{i[k_i-1]}$, respectively, using Lemma 2.

5. An Example

We only considered the Ξ_{CON} assignment problem where $\varphi = 15, \aleph = 3, \varphi_1 = \varphi_2 = \varphi_3 = 5, \ell = 2, \gamma = 4, \widetilde{s}_1 = 2, \widetilde{s}_2 = 3, \widetilde{s}_3 = 1$; the parameters of job J_{ih} ($i = 1, 2, 3; h = 1, \dots, 5$) are given in Table 2, α_{ih} and β_{ih} of \widetilde{PDW} ($i = 1, 2, 3; j = 1, \dots, 5$) are presented in Table 3.

Table 2. Job parameters.

\widetilde{G}_i	\widetilde{G}_1					\widetilde{G}_2					\widetilde{G}_3				
J_{ih}	J_{11}	J_{12}	J_{13}	J_{14}	J_{15}	J_{21}	J_{22}	J_{23}	J_{24}	J_{25}	J_{31}	J_{32}	J_{33}	J_{34}	J_{35}
a_{ih}	14	15	13	16	11	13	16	17	18	17	16	19	23	21	18
b_{ih}	2	3	3	1	2	2	3	4	5	3	4	4	5	6	3
\bar{u}_{ih}	4	3	4	10	5	6	5	4	3	5	3	4	4	3	5
θ_{ih}	15	13	9	10	17	15	11	19	20	9	16	15	21	15	18
g_{ih}	4	5	6	2	7	3	6	5	3	4	5	7	8	9	11

Table 3. Position-dependent weights.

\widetilde{G}_i	\widetilde{G}_1				\widetilde{G}_2				\widetilde{G}_3						
α_{ih}	6	4	8	6	10	8	10	9	10	13	9	14	10	16	11
β_{ih}	10	11	14	8	12	13	12	15	11	14	16	17	11	18	12

For problem

$$1 \left| p_{ih} = a_{ih} - b_{ih}u_{ih}, \Xi_{CON,GT,PDW} \left| \sum_{i=1}^N \sum_{h=1}^{\varphi_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i), \right. \right.$$

from Algorithm 1 and Lemma 2, $8 + 12 - (6 + 4 + 8) = 2 \leq 4 \times 5 \leq 14 + 8 + 12 - (6 + 4) = 24$; hence $k_1 = 3$. Similarly, $k_2 = 3, k_3 = 2$. Values $\vartheta_{jh}^{[1]}$ are given in Table 4 when \widetilde{G}_1 was scheduled at the r th position. Table 4 shows that the optimal job sequence was $\varphi_{11}^* = \{J_{11} \rightarrow J_{13} \rightarrow J_{15} \rightarrow J_{12} \rightarrow J_{14}\}$, similarly, $\varphi_{12}^* = \{J_{11} \rightarrow J_{13} \rightarrow J_{15} \rightarrow J_{12} \rightarrow J_{14}\}$, $\varphi_{13}^* = \{J_{11} \rightarrow J_{13} \rightarrow J_{15} \rightarrow J_{12} \rightarrow J_{14}\}$; For group \widetilde{G}_2 , we have $\varphi_{21}^* = \{J_{25} \rightarrow J_{21} \rightarrow J_{22} \rightarrow J_{23} \rightarrow J_{24}\}$, $\varphi_{22}^* = \{J_{25} \rightarrow J_{21} \rightarrow J_{22} \rightarrow J_{23} \rightarrow J_{24}\}$, $\varphi_{23}^* = \{J_{25} \rightarrow J_{21} \rightarrow J_{22} \rightarrow J_{23} \rightarrow J_{24}\}$; For group \widetilde{G}_3 , we have $\varphi_{31}^* = \{J_{32} \rightarrow J_{33} \rightarrow J_{34} \rightarrow J_{35} \rightarrow J_{31}\}$, $\varphi_{32}^* = \{J_{32} \rightarrow J_{33} \rightarrow J_{34} \rightarrow J_{35} \rightarrow J_{31}\}$, $\varphi_{33}^* = \{J_{32} \rightarrow J_{33} \rightarrow J_{34} \rightarrow J_{35} \rightarrow J_{31}\}$.

According to Step 4 of Algorithm 1, the values of χ_{ir} are given in Table 5. From Table 5, we have $q^* = \{\widetilde{G}_2 \rightarrow \widetilde{G}_3 \rightarrow \widetilde{G}_1\}$. The optimal jobs sequences are $\varphi_{21}^* = \{J_{25} \rightarrow J_{21} \rightarrow J_{22} \rightarrow J_{23} \rightarrow J_{24}\}$, $\varphi_{32}^* = \{J_{32} \rightarrow J_{33} \rightarrow J_{34} \rightarrow J_{35} \rightarrow J_{31}\}$, and $\varphi_{13}^* = \{J_{11} \rightarrow J_{13} \rightarrow J_{15} \rightarrow J_{12} \rightarrow J_{14}\}$. The optimal resource allocations corresponding to the sequence are $u_{25}^* = \bar{u}_{25} = 5, u_{21}^* = \bar{u}_{21} = 6, u_{22}^* = \bar{u}_{22} = 5, u_{23}^* = \bar{u}_{23} = 4, u_{24}^* = \bar{u}_{24} = 3, u_{32}^* = \bar{u}_{32} = 4, u_{33}^* = \bar{u}_{33} = 4, u_{34}^* = \bar{u}_{34} = 3, u_{35}^* = \bar{u}_{35} = 5, u_{31}^* = \bar{u}_{31} = 3, u_{11}^* = \bar{u}_{11} = 4, u_{13}^* = \bar{u}_{13} = 4, u_{15}^* = \bar{u}_{15} = 5, u_{12}^* = \bar{u}_{12} = 3, u_{14}^* = \bar{u}_{14} = 10$. The optimal due-dates are $d_1^* = C_{22} = 7, d_2^* = C_{34} = 18$, and $d_3^* = C_{13} = 38$.

Table 4. Values $\vartheta_{jh}^{[1]}$ for \widetilde{G}_1 (bold numbers are the optimal solution).

	$h = 1$	$h = 2$	$h = 3$	$h = 4$	$h = 5$
J_{11}	376	412	436	376	328
J_{12}	375	411	435	375	327
J_{13}	84	90	94	84	76
J_{14}	380	416	440	380	332
J_{15}	95	101	105	95	87

Table 5. Values χ_{ir} of the linear problem (bold numbers are the optimal solution).

	$r = 1$	$r = 2$	$r = 3$
\widetilde{G}_1	1398	958	518
\widetilde{G}_2	758	538	318
\widetilde{G}_3	1265	925	585

Similarly, for problem

$$1 \left| p_{ih} = \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell, \Xi_{CON,GT,PDW} \left| \sum_{i=1}^N \sum_{h=1}^{\varphi_i} (\alpha_{ih}E_{i[h]} + \beta_{ih}T_{i[h]} + g_{ih}u_{ih} + \gamma d_i), \right. \right.$$

$k_1 = 3, k_2 = 3, k_3 = 2$. According to Lemma 7, for $i = 1$, the optimal job sequence is $\varphi_1^* = \{J_{11} \rightarrow J_{13} \rightarrow J_{14} \rightarrow J_{12} \rightarrow J_{15}\}$, for $i = 2$, the optimal sequence is $\varphi_2^* = \{J_{22} \rightarrow J_{21} \rightarrow J_{25} \rightarrow J_{24} \rightarrow J_{23}\}$, for $i = 3$, the optimal sequence is $\varphi_3^* = \{J_{33} \rightarrow J_{34} \rightarrow J_{31} \rightarrow J_{32} \rightarrow J_{35}\}$. According to Equation (40), the values of χ_{ir} are shown in Table 6, where

we have $q^* = \{\widetilde{G}_1, \widetilde{G}_2, \widetilde{G}_3\}$. The optimal values of resource allocation corresponding to the sequence are $u_{11}^* = 18.8988, u_{13}^* = 12.1237, u_{14}^* = 19.1293, u_{12}^* = 15.9477, u_{15}^* = 16.2534, u_{22}^* = 11.7285, u_{21}^* = 19.3098, u_{25}^* = 13.2931, u_{24}^* = 22.8943, u_{23}^* = 16.9961, u_{33}^* = 13.0158, u_{34}^* = 11.3185, u_{31}^* = 16.1322, u_{32}^* = 12.4474, u_{35}^* = 26.3904$. The optimal due dates are $d_1^* = C_{14} = 3.4544, d_2^* = C_{25} = 10.0560$, and $d_3^* = C_{34} = 17.4282$.

Table 6. Values χ_{ir} of the convex problem (bold numbers are the optimal solution).

	$r = 1$	$r = 2$	$r = 3$
\widetilde{G}_1	690.1632	576.7474	492.4878
\widetilde{G}_2	751.6441	622.7040	465.1836
\widetilde{G}_3	1043.3726	910.8574	724.1781

6. Conclusions

In this article, we investigated single-machine group technology scheduling with \widetilde{CPT} . Under Ξ_{CON}/Ξ_{SLK} assignments, the goal is to find the job and group sequences, resource allocation, and due-date assignment, such that the sum of scheduling and resource-allocation costs is minimized. For $\wp_i = \widehat{N}$ ($i = 1, \dots, \aleph$), we demonstrated that this problem is polynomially solvable. Future work could explore job (or flow) shop problems (see Guo et al. [33], and Karacan et al. [34]) with group technology and controllable processing times to study the general version of

$$1 \left| p_{ih}, \Xi_{CON}/\Xi_{SLK}, GT, \widehat{PDW} \right| \sum_{i=1}^{\aleph} \sum_{h=1}^{\wp_i} (\alpha_{ih} E_{i[h]} + \beta_{ih} T_{i[h]} + g_{ih} u_{ih} + \gamma d_i / q_i),$$

where $p_{ih} \in \left\{ a_{ih} - b_{ih} u_{ih}, \left(\frac{\theta_{ih}}{u_{ih}} \right)^\ell \right\}$ (e.g., the proposed cuckoo search algorithm of Xie et al. [35]). Future work could also consider problems regarding maintenance activity (see Wu et al. [36]).

Author Contributions: Conceptualization, W.L. and X.W.; methodology, W.L. and X.W.; writing, review and editing, W.L. and X.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the National Natural Science Regional Foundation of China (72061029 and 71861031).

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Lu, Y.-Y.; Li, G.; Wu, Y.-B.; Ji, P. Optimal due-date assignment problem with learning effect and resource-dependent processing times. *Optim. Lett.* **2014**, *8*, 113–127. [\[CrossRef\]](#)
- Liu, L.; Wang, J.-J.; Wang, X.-Y. Single machine due-window assignment scheduling with resource-dependent processing times to minimise total resource consumption cost. *Int. J. Prod. Res.* **2016**, *54*, 1186–1195. [\[CrossRef\]](#)
- Li, L.; Wang, J.-J. Scheduling jobs with deterioration effect and controllable processing time. *Neural Comput. Appl.* **2018**, *29*, 1163–1170. [\[CrossRef\]](#)
- Lu, Y.-Y.; Liu, J.-Y. A note on resource allocation scheduling with position-dependent workloads. *Eng. Optim.* **2018**, *50*, 1810–1827. [\[CrossRef\]](#)
- Geng, X.-N.; Wang, J.-B.; Bai, D. Common due date assignment scheduling for a no-wait flowshop with convex resource allocation and learning effect. *Eng. Optim.* **2019**, *51*, 1301–1323. [\[CrossRef\]](#)
- Sun, X.; Geng, X.-N.; Wang, J.-B.; Liu, F. Convex resource allocation scheduling in the no-wait flowshop with common flow allowance and learning effect. *Int. J. Prod. Res.* **2019**, *57*, 1873–1891. [\[CrossRef\]](#)
- Liu, W.-W.; Jiang, C. Flow shop resource allocation scheduling with due date assignment, learning effect and position-dependent weights. *Asia-Pac. J. Oper. Res.* **2020**, *37*, 2050014. [\[CrossRef\]](#)
- Lu, Y.-Y.; Wang, T.-T.; Wang, R.-Q.; Li, Y. A note on due-date assignment scheduling with job-dependent learning effects and convex resource allocation. *Eng. Optim.* **2021**, *53*, 1273–1281. [\[CrossRef\]](#)

9. Zhao, S. Resource allocation flowshop scheduling with learning effect and slack due window assignment. *J. Ind. Manag. Optim.* **2021**, *17*, 2817–2835. [[CrossRef](#)]
10. Lv, D.-Y.; Wang, J.-B. Study on resource-dependent no-wait flow shop scheduling with different due-window assignment and learning effects. *Asia-Pac. J. Oper. Res.* **2021**, *38*, 2150008. [[CrossRef](#)]
11. Tian, Y. Single-machine due-window assignment scheduling with resource allocation and generalized earliness/tardiness penalties. *Asia-Pac. J. Oper. Res.* **2022**, *39*, 2150041. [[CrossRef](#)]
12. Wang, Y.-C.; Wang, S.-H.; Wang, J.-B. Resource allocation scheduling with position-dependent weights and generalized earliness-tardiness cost. *Mathematics* **2023**, *11*, 222. [[CrossRef](#)]
13. Liu, S.-C. Common due-window assignment and group scheduling with position-dependent processing times. *Asia-Pac. J. Oper. Res.* **2015**, *32*, 1550045. [[CrossRef](#)]
14. Wang, J.-B.; Liu, L.; Wang, J.-J.; Li, L. Makespan minimization scheduling with ready times, group technology and shortening job processing times. *Comput. J.* **2018**, *61*, 1422–1428. [[CrossRef](#)]
15. Huang, X. Bicriterion scheduling with group technology and deterioration effect. *J. Appl. Math. Comput.* **2019**, *60*, 455–464. [[CrossRef](#)]
16. Liu, F.; Yang, J.; Lu, Y.-Y. Solution algorithms for single-machine group scheduling with ready times and deteriorating jobs. *Eng. Optim.* **2019**, *51*, 862–874. [[CrossRef](#)]
17. Wang, L.-Y.; Liu, M.; Wang, J.-B.; Lu, Y.-Y.; Liu, W.-W. Optimization for due-date assignment single-machine scheduling under group technology. *Complexity* **2021**, *2021*, 6656261. [[CrossRef](#)]
18. Wang, J.-B.; Jia, X.; Yan, J.-X.; Wang, S.-H.; Qian, J. Single machine group scheduling problem with makespan objective and a proportional linear shortening. *RAIRO-Oper. Res.* **2022**, *56*, 1523–1532. [[CrossRef](#)]
19. Chen, K.; Han, S.; Huang, H.; Ji, M. A group-dependent due window assignment scheduling problem with controllable learning effect. *Asia-Pac. J. Oper. Res.* **2023**, *40*, 2250025. [[CrossRef](#)]
20. Shabtay, D.; Itskovich, Y.; Yedidsion, L.; Oron, D. Optimal due date assignment and resource allocation in a group technology scheduling environment. *Comput. Oper. Res.* **2010**, *37*, 2218–2228. [[CrossRef](#)]
21. Zhu, Z.; Sun, L.; Chu, F.; Liu, M. Single-machine group scheduling with resource allocation and learning effect. *Comput. Ind. Eng.* **2011**, *60*, 148–157. [[CrossRef](#)]
22. Wang, D.; Huo, Y.; Ji, P. Single-machine group scheduling with deteriorating jobs and allotted resource. *Optim. Lett.* **2014**, *8*, 591–605. [[CrossRef](#)]
23. Wang, J.-B.; Liang, X.-X. Group scheduling with deteriorating jobs and allotted resource under limited resource availability constraint. *Eng. Optim.* **2019**, *51*, 231–246. [[CrossRef](#)]
24. Liang, X.-X.; Liu, M.; Feng, Y.-B.; Wang, J.-B.; Wen, L.-S. Solution algorithms for single-machine resource allocation scheduling with deteriorating jobs and group technology. *Eng. Optim.* **2020**, *52*, 1184–1197. [[CrossRef](#)]
25. Yan, J.-X.; Ren, N.; Bei, H.-B.; Bao, H.; Wang, J.-B. Study on resource allocation scheduling problem with learning factors and group technology. *J. Ind. Manag. Optim.* **2023**, *19*, 3419–3435. [[CrossRef](#)]
26. Chen, Y.; Ma, X.; Zhang, F.; Cheng, Y. On optimal due date assignment without restriction and resource allocation in group technology scheduling. *J. Comb. Optim.* **2023**, *45*, 64. [[CrossRef](#)]
27. Gordon, V.; Proth, J.M.; Chu, C.B. A survey of the state-of-the-art of common due date assignment and scheduling research. *Eur. J. Oper.* **2002**, *139*, 1–25. [[CrossRef](#)]
28. Gordon, V.; Proth, J.M.; Chu, C.B. Due date assignment and scheduling: SLK, TWK and other due date assignment models. *Prod. Plan. Control.* **2002**, *13*, 117–132. [[CrossRef](#)]
29. Liu, W.; Hu, X.; Wang, X. Single machine scheduling with slack due dates assignment. *Eng. Optim.* **2017**, *49*, 709–717. [[CrossRef](#)]
30. Wang, X.; Liu, W.; Li, L.; Zhao, P.; Zhang, R. Due date assignment scheduling with positional-dependent weights and proportional setup times. *Math. Biosci. Eng.* **2022**, *19*, 5104–5119. [[CrossRef](#)]
31. Wang, J.-B.; Wang, S.-H.; Gao, K.; Liu, M.; Jia, X. Due-window assignment methods and scheduling with generalized positional-dependent weights. *Asia-Pac. J. Oper. Res.* **2022**, *39*, 2250028. [[CrossRef](#)]
32. Hardy, G.H.; Littlewood, J.E.; Polya, G. *Inequalities*, 2nd ed.; Cambridge University Press: Cambridge, UK, 1967. [[CrossRef](#)]
33. Guo, J.; Luo, Q.; Liang, P.; Ouyang, J. A GAPN approach for the flexible job-shop scheduling problem with indirect energy and time-of-use electricity pricing. *Processes* **2022**, *10*, 832.
34. Karacan, I.; Senvar, O.; Bulkan, S. A novel parallel simulated annealing methodology to solve the no-wait flow shop scheduling problem with earliness and tardiness objectives. *Processes* **2023**, *11*, 454. [[CrossRef](#)]
35. Xie, X.; Zheng, Y.; Mu, T.; Wan, F.; Dong, H. Solving the two-crane scheduling problem in the pre-steelmaking process. *Processes* **2023**, *11*, 549. [[CrossRef](#)]
36. Wu, W.; Lv, D.-Y.; Wang, J.-B. Two due-date assignment scheduling with location-dependent weights and a deteriorating maintenance activity. *Systems* **2023**, *11*, 150. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.