



Systematic Review

A Systematic Review of Ebstein's Anomaly with Left Ventricular Noncompaction

Suma K. Thareja^{1,2}, Michele A. Frommelt^{3,4}, Joy Lincoln^{3,4} , John W. Lough¹ , Michael E. Mitchell^{2,3,4} and Aoy Tomita-Mitchell^{2,3,4,*}

- ¹ Department of Cell Biology, Neurobiology, and Anatomy, Medical College of Wisconsin, Milwaukee, WI 53226, USA; sthareja@mcw.edu (S.K.T.); jlough@mcw.edu (J.W.L.)
² Department of Surgery, Division of Congenital Heart Surgery, Medical College of Wisconsin, Milwaukee, WI 53226, USA; memitchell@mcw.edu
³ Department of Pediatrics, Division of Pediatric Cardiology, Children's Wisconsin, Milwaukee, WI 53226, USA; mafrom@mcw.edu (M.A.F.); jlincoln@mcw.edu (J.L.)
⁴ Herma Heart Institute, Children's Wisconsin, Milwaukee, WI 53226, USA
* Correspondence: amitchell@mcw.edu

Abstract: Traditional definitions of Ebstein's anomaly (EA) and left ventricular noncompaction (LVNC), two rare congenital heart defects (CHDs), confine disease to either the right or left heart, respectively. Around 15–29% of patients with EA, which has a prevalence of 1 in 20,000 live births, commonly manifest with LVNC. While individual EA or LVNC literature is extensive, relatively little discussion is devoted to the joint appearance of EA and LVNC (EA/LVNC), which poses a higher risk of poor clinical outcomes. We queried PubMed, Medline, and Web of Science for all peer-reviewed publications from inception to February 2022 that discuss EA/LVNC and found 58 unique articles written in English. Here, we summarize and extrapolate commonalities in clinical and genetic understanding of EA/LVNC to date. We additionally postulate involvement of shared developmental pathways that may lead to this combined disease. Anatomical variation in EA/LVNC encompasses characteristics of both CHDs, including tricuspid valve displacement, right heart dilatation, and left ventricular trabeculation, and dictates clinical presentation in both age and severity. Disease treatment is non-specific, ranging from symptomatic management to invasive surgery. Apart from a few variant associations, mainly in sarcomeric genes *MYH7* and *TPM1*, the genetic etiology and pathogenesis of EA/LVNC remain largely unknown.

Keywords: Ebstein's anomaly; left ventricular noncompaction; congenital heart disease; cardiomyopathy; genetic etiology



Citation: Thareja, S.K.; Frommelt, M.A.; Lincoln, J.; Lough, J.W.; Mitchell, M.E.; Tomita-Mitchell, A. A Systematic Review of Ebstein's Anomaly with Left Ventricular Noncompaction. *J. Cardiovasc. Dev. Dis.* **2022**, *9*, 115. <https://doi.org/10.3390/jcdd9040115>

Academic Editor: Jau-Nian Chen

Received: 12 March 2022

Accepted: 7 April 2022

Published: 13 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Congenital heart defects (CHDs) are cardiogenic malformations present at or before birth that account for a high burden of neonatal morbidity and mortality in the developed world. As improvements in cardiac medicine enable patients with CHDs to live longer, there remains an unmet need for tailored medical care and treatment. Here, we provide a brief overview of two rare CHDs—Ebstein's anomaly (EA) and left ventricular noncompaction (LVNC)—and thoroughly review publications that discuss combined EA/LVNC.

1.1. Ebstein's Anomaly (EA)

Reported EA prevalence ranges from one in 200,000 to, more commonly, one in 20,000 live births, and accounts for less than one percent of CHD cases [1–4]. Improved patient health outcomes, advanced diagnostic imaging, and genomic technology likely contribute to this epidemiological shift [4].

EA mainly affects the tricuspid valve and right ventricle. Specific EA characteristics include the attachment of posterior and septal leaflets to the ventricular septum, apical

displacement of the valve, redundant and fenestrated anterior leaflet, atrialization of the portion of right ventricle above the valve, and dilation of the right ventricle and atrioventricular junction [1,5,6].

Significant EA pathology originates from the right heart dilation, and consequently, over two-thirds of EA patients display enlargement of the right heart, including the atrialized inlet portion of the right ventricle and often the functional right ventricular apex and outflow tract [1]. Pronounced dilation also narrows the left ventricular chamber (visibly shaped as a crescent on an echocardiogram, or ECHO) and may lead to heart failure from left ventricular outflow tract obstruction [1,7].

Clinical heterogeneity of EA patients ranges in anatomic and physiologic severity, and as such, treatment from medication to valve repair is highly individualized [8,9]. Fetal presentation or severity of clinical symptoms increases risk of mortality [10]. Common EA-associated cardiac comorbidities are atrial septal defect (ASD), Wolff–Parkinson–White (WPW) syndrome, and right ventricular outflow tract obstruction [11]. Neonates born on the verge of viability commonly exhibit symptoms of heart failure (e.g., poor feeding, failure to thrive, and dyspnea) [1]. Survival to adulthood is variable and dependent on clinical severity; thus, EA prevalence in adults is not thoroughly enumerated.

Categorization of EA as mild, moderate, or severe is based on varied classification systems: (1) Carpentier et al. [12] assess freedom of valve leaflet movement and severity of outflow tract obstruction; (2) Celermajer et al. [10] grade severity based on volumetric chamber measurements from ECHO [10]; and Dearani et al. [13] review both the cardiac morphology on ECHO and leaflet anatomy from surgical observation.

1.2. Left Ventricular Noncompaction (LVNC)

EA also commonly manifests with left ventricular noncompaction (LVNC), a cardiomyopathy of ventricular myocardium that does not transit from a weak and “spongy” stage to a compacted state during development [14,15]. Morphological characteristics include abnormal trabeculations and deep intra-trabecular recesses adjacent to the endothelium that resemble the morphology of the right ventricle.

Common LVNC-associated cardiac comorbidities include ventricular septal defect (VSD), ASD, and patent ductus arteriosus (PDA). One pathologic predictor of LVNC is a greater base–apex strain, resulting from longitudinal deformation [16].

Prevalence of LVNC is even more variable than EA, and diagnosis may be missed in patients with asymptomatic clinical presentation or absent imaging. Similar to EA, early LVNC presentation in infancy increases the risk of mortality [17]. Often, the left ventricle is dilated, or hypertrophic, and severe cases may result in heart failure from either or both systolic and diastolic dysfunction [15]. Men are more likely to present with noncompaction (56–82% of cases) than women [18].

At least eight different LVNC phenotypes that range from benign to restrictive pathology with variable clinical associations such as ventricular dilation, arrhythmias, and systolic dysfunction exist [15]. Three common criteria for a diagnosis of LVNC include a ratio of the length of the compacted wall (x) to the trabeculated wall (y) being less than 0.5 (i.e., $x/y < 0.5$) on short-axis ECHO [19], a noncompacted to compacted ratio (NC:C) greater than 1.4 in children or 2 in adults, also on short-axis ECHO [20], and a NC:C ratio greater than 2.3 on long-axis cardiac MRI [21].

1.3. Ebstein’s Anomaly with Left Ventricular Noncompaction (EA/LVNC)

Illustrated in Figure 1 are the morphological alterations of combined EA and LVNC (EA/LVNC), which is a specific category of EA patients that was first described by Attenhofer Jost et al. in 2004 [22]. The spectrum of phenotypes for both CHDs underscores both the putative genetic etiology and the complexity of the $G \times E$ interaction during disease development.




Figure 1. Illustration of EA/LVNC morphology: (A) normal heart; (B) morphological defects seen in EA/LVNC. Common characteristics depicted here are the apical displacement of the tricuspid valve, dilation of the right atrioventricular junction, trabeculated left ventricular myocardium, and an atrial septal defect (a common comorbidity of EA/LVNC). EA/LVNC; Ebstein’s anomaly with left ventricular noncompaction.

2. Methods

We followed the systematic review guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [23,24] and identified all peer-reviewed publications from inception to February 2022 that considered EA/LVNC (Figure 2). We accessed PubMed, Medline, and Web of Science with the search terms “(Ebstein’s anomaly) and (left ventricular noncompaction)”, “(Ebstein’s anomaly) AND (left ventricular noncompaction)”, “(Ebstein’s anomaly) AND (LVNC)”, “(Ebstein’s anomaly) AND (left ventricular hypertrabeculation)”, “(Ebstein’s anomaly) AND (noncompaction)”, and “(Ebstein’s malformation) AND (noncompaction)” for relevant articles. We also queried clinical trials or studies noted on ClinicalTrials.gov and clinicaltrialsregister.eu. We then tabulated and created tables of EA/LVNC clinical signs, imaging modalities, treatments, clinical trials, and genetic associations. To eliminate risk of bias while S.K.T. independently reviewed all articles, we only excluded publications that were not written in English or did not consider combined EA/LVNC.




Figure 2. PRISMA 2020 flow diagram [23,24] for this systematic review on EA/LVNC.

3. Results

3.1. We Identified 58 Peer-Reviewed Publications That Discuss EA/LVNC

As presented in the PRISMA flow diagram (Figure 2) [23,24], we identified a total of 358 papers from inception to February 2022 from the use of the search terms applied to manuscripts. Of these, 287 were redundant. Among the 71 unique articles, we excluded 6 following title/abstract screening [25–30]. We retrieved 65 full-text articles and further excluded 2 articles that were not written in English [31,32] and 5 with no direct focus on EA/LVNC in association with each other or on a morphologically similar spectrum [1,5,6,33,34]. This resulted in 58 unique articles published between 2004 and 2022, including sporadic case reports [22,35–67], familial case reports [68–75], large-scale clinical or genetic studies [7,17,18,76–83], and reviews on related clinical manifestations [84–88].

3.2. EA/LVNC Patients Present with Clinical Signs of Both CHDs

Dual EA/LVNC patients present with clinical characteristics of both myocardial disorders (Figure 1B) [22,76,77]. As EA commonly manifests with LVNC, the incidence of EA with LVNC is estimated to be between 15% and 29% [78,79,81,82] and much lower (<5%) when only adult patients are considered [80]. There appears to be no clear increased prevalence, risk, or differing phenotypes of EA or LVNC associated with race.

Table 1 summarizes common clinical signs of EA/LVNC based on age. Childhood presentation is associated with signs of heart failure, while late adulthood is associated with asymptomatic or incidental finding. All other age ranges in case reports display variable clinical signs, including exertional dyspnea, chest pain, and arrhythmia. Among all sporadic and familial case reports, two most common clinical signs are exertional dyspnea and systolic murmur from tricuspid regurgitation (Table 2). Larger-scale studies of EA, LVNC, or EA/LVNC are outlined in Table 3.

Table 1. Common clinical signs based on age of sporadic EA/LVNC case presentation.

Age of Sporadic Case Presentation	Common Clinical Signs
Prenatal/Infancy (<1 year) [40,42,47,56,57,60,63]	Cyanosis
Childhood (1–12 years) [41,50]	Heart Failure Systolic Murmur
Adolescence (13–20 years) [22,39,43,44,49,54,59,67] OR Early Adulthood (21–40 years) [22,37,44–46,51–53,58,61] OR Middle Adulthood (41–60 years) [35,36,38,48,55,62,64,66]	Progressive or Exertional Dyspnea Systolic Murmur Palpitations or Tachycardia Chest Pain or Discomfort Arrhythmia
Late Adulthood (61 years+) [65]	Progressive or Exertional Dyspnea Asymptomatic/Incidental Finding

Patients born with both EA and LVNC and diagnosed in infancy have a higher risk of poor outcomes such as heart failure and sudden cardiac death and manifest with anatomical variations that pose an additional surgical challenge [7,55,82,83]. Thus, a decreased incidence of dual EA/LVNC presentation in adulthood appears to correlate with increased morbidity. Similar to individual EA or LVNC, septal defects (ASD or VSD) are the most common comorbidities of EA/LVNC [22,44,47,63,69,70].

In cases of combined EA/LVNC, typically one of the two conditions is present in the relatively mild state. Some case reports describe patients initially diagnosed by EA and a later diagnosis of LVNC following an ischemic attack or unexplained clinical phenomena that prompted additional imaging [38,62,66]. Examples also exist where a patient will exhibit EA, LVNC, and other cardiac abnormalities [36,52], yet typically one of the conditions presents latently or only intermittently. In certain cases, significant remodeling of the heart is noted, presumably from acclimatization to an anatomical and hence physiological challenge [51]. The converse conclusion may be that in most instances where morphological features of both CHDs appear—or would have appeared if the patient survived—the ontological pathways that react to the dual challenge were not available or failed. Clinicians and researchers studying EA/LVNC postulate that there may be some link between both diseases as evidenced by a strong, if not complete, genetic component, and thus, an abnormal pathology would presumably exist in those affected from birth.

3.3. EA/LVNC Patients Commonly Present with Arrhythmia

A clinical association of arrhythmia for patients with EA/LVNC originated as early as 1978. Monibi et al. evaluated 17 newborns with EA and found that 13 presented with high normal or elevated hematocrit without cyanosis, and 9 of 13 who had LV catheterization and cineangiography displayed abnormal contraction with unusually elongated and distorted long axis and irregularity of septal and lateral walls [89]. Of the 13 patients, 12 had abnormal contraction, hypokinesis, or dyskinesis [89].

Table 2. Clinical signs, imaging modalities, and interventions noted among all EA/LVNC case reports.

Clinical Signs	Imaging Modalities	Interventions/Therapies
		Mechanical Ventilation [42,50,57]
	2-D ECHO [7,22,35–59,61–75]	Radiofrequency Ablation [35,58,62]
Arrhythmia [35,37,43,45,51,55,61,63,65,73,74]	3-D ECHO [36,39,46,49,53,61,65,75]	Arterioplasty/Annuloplasty [22]
Asymptomatic/Incidental Finding [36,65,66,75]	Color Flow Doppler ECHO [36,37,44,46–48,51,54,58,59,61,63,68–70,73,75]	Other Cardiac Surgical Repair [7,41,47,53,56,70]
Chest Pain or Discomfort [46,58,61,62]	Cardiac or Coronary CT Angiography [35,46,56,62]	Cardiac Transplantation [22,44,50,69]
Cyanosis [47,57,74]	Cardiac MRI [37,38,40,41,45,50,51,53,55,56,62,64–66,69,73]	Atrial Septal Occluder (PFO Closure) [38]
Exertional Dyspnea [22,35,38,41–45,49,51,53,55,64,65,68,69,74]	Catheterization [53,73]	LVAD/BiVAD [43,53,74]
Heart Failure [50,73,74]	Chest Radiograph [7,38,43,50,51,57,58,61]	Pacemaker/ICD [35,55,73,74]
Palpitations or Tachycardia [35,41,44,45,58,62]	ECG [35–37,41,43,44,47–49,51,52,54,58,61,63,65,73–75]	IABP [53]
Systolic Murmur [39,40,45,48,49,54,59,66,67]	Electrophysiological Study [35,52,73,74]	Inotropic Support (Epinephrine/Milirinone) [53]
		Anticoagulation (Aspirin/Warfarin/Heparin) [35,39,43–45,49,52,59,65,73]
		Heart Failure and Anti-arrhythmic Medications (Beta-Blockers/ACE inhibitors/Diuretics/Digoxin) [43–47,51,52,57,65,73,74]

BiVAD: Biventricular Assist Device, CT: Computerized Tomography, ECG: Electrocardiogram, ECHO: Echocardiography, IABP: Intra-aortic Balloon Pump, ICD: Implantable Cardioverter-Defibrillator, LVAD: Left Ventricular Assist Device, MRI: Magnetic Resonance Imaging, PFO: Patent Foramen Ovale.

Two decades hence, cineangiography of 26 adults with EA identified that 16 presented with abnormal LV contraction (hypokinesia in 10, dyskinesia in 6, and premature diastolic distention of the anterobasal wall in 8) [90].

Table 3. Clinical Studies that discuss EA/LVNC.

Study	Published	Focus	Center	Subjects	Gender	EA/LVNC	Results
Attenhofer Jost et al. [78]	2005	Clinical study of EA	Mayo Clinic Rochester, MN, USA	106 patients aged 0–52 years.	39.7% M 60.3% F	<ul style="list-style-type: none"> 19 EA patients with LVNC (18%) 	<ul style="list-style-type: none"> 39% of EA patients exhibited LV abnormalities including LVNC, bicuspid aortic valve, VSD, and mitral valve abnormality Decreased trend of the ECG QRS axis mean in EA/LVNC (12°) compared with all patients (36°)

Table 3. Cont.

Study	Published	Focus	Center	Subjects	Gender	EA/LVNC	Results
El-Menyar et al. [18]	2007	Clinical study of LVNC in Qatar	Hamad General Hospital Doha, Qatar	12 patients aged 0–37 years.	33% M 67% F	<ul style="list-style-type: none"> 1 LVNC patient with EA 	<ul style="list-style-type: none"> 25% mortality rate 50% LVNC patients associated with CHD (VSD, PS, EA, coarctation) Poor prognosis associated with biventricular noncompaction
Reemtsen et al. [7]	2007	Clinical study of EA	Children’s Hospital Los Angeles Los Angeles, CA, USA	12 patients aged 0–17 years.	58% M 42% F	<ul style="list-style-type: none"> All neonates with EA had severe LV dysfunction 	<ul style="list-style-type: none"> 12 of 16 patients aged 0–17 years. survived neonatal RV exclusion surgery 11 of 12 survivors received a bidirectional Glenn shunt and 6 were completely palliated with a Fontan Prior to Glenn and Fontan, patients had a decreased CT ratio, Great Ormond Street ratio, RV/LV ratio, septal impingement from a/b approaching ratio, and shortening fraction; all these parameters improved following Glenn and Fontan
Tsai et al. [17]	2009	Clinical study of LVNC	Riley Hospital for Children Indianapolis, IN, USA	46 patients	50% M 50% F	<ul style="list-style-type: none"> 5 LVNC patients with EA 	<ul style="list-style-type: none"> 54% heart failure 52% decreased LV ejection fraction 78% associated cardiac defects (ASD, VSD, PDA, EA) 80% ECG abnormalities 20% mortality rate not correlated to ejection fraction, morphological defect, or arrhythmia
Stähli et al. [79]	2013	Clinical study of LVNC	University Children’s Hospital Zurich, Switzerland	202 patients	79% M 21% F	<ul style="list-style-type: none"> 6 EA patients with LVNC (15%) 	<ul style="list-style-type: none"> 24% associated cardiac defects (aortic valve abnormalities, EA, TOF, DORV)
Pignatelli et al. [82]	2014	Clinical study of EA or EA/LVNC	Texas Children’s Hospital Houston, TX, USA	61 infants		<ul style="list-style-type: none"> 51 infants with EA 10 EA infants with LVNC (16.4%) 	<ul style="list-style-type: none"> EA/LVNC cohort trended in earlier detection (9/10 patients were diagnosed at birth) and a higher risk of adverse outcomes (progressive LV dysfunction) than patients with EA alone

Table 3. Cont.

Study	Published	Focus	Center	Subjects	Gender	EA/LVNC	Results
Kumor et al. [80]	2018	Clinical study of EA	Institute of Cardiology Warsaw, Poland	84 patients aged 16–71 years.	41% M 59% F	<ul style="list-style-type: none"> 4 EA patients with LVNC (4.8%) 	<ul style="list-style-type: none"> ASD type II (27.3%) and WPW syndrome (10.7%) were common in EA patients EA/LVNC patients alone in this cohort suffered cardiac arrest or ventricular arrhythmia
Hirono et al. [83]	2020	Clinical and genetic study of LVNC	University of Toyama Toyama, Japan;	53 Japanese probands aged 1–14 years.	47% M 53% F	<ul style="list-style-type: none"> 7 LVNC patients with EA 	<ul style="list-style-type: none"> LVNC/CHD patients had lower ejection fractions, thickened trabeculations, and worse prognosis when compared with age-matched patients with ventricular septal defects Heart failure, LV ejection fraction of <24%, LV end-diastolic diameter z-score of >8.56, and the LV NC:C ratio >8.33 at the last visit were risk factors for survival 30 genetic variants in 28 patients with LVNC and CHD in the genes (50% sarcomeric): MYH7, TPM1, ACTC1, ANK2, COL4A1, DAAM1, DSG2, DSP, FGF16, FGFR2, HCN4, JUP, MYBPC3, MYH6, MYL2, PKP2, PRDM16, RYR2, and TAZ
Marques et al. [81]	2020	CHD study of postnatal heart specimens	Heart Institute (InCor), University of Sao Paulo Medical School, Brazil	259 postnatal hearts with 87.3% aged less than 18 years.	49% M 51% F	<ul style="list-style-type: none"> 23 patients with EA 	<ul style="list-style-type: none"> Prevalence of LVNC in EA patients was either 28.6%, 9.5%, or 0% based on LVNC diagnostic criteria using Chin’s, [91] Jenni’s [20] or Petersen’s [21] approaches VSD hearts presented significantly higher with LVNC regardless of the three methods of LVNC diagnosis

CHD: Congenital Heart Defect, DORV: Double Outlet Right Ventricle, EA: Ebstein’s Anomaly, ECG: Electrocardiogram, LV: Left Ventricle, LVNC: Left Ventricular Noncompaction, PDA: Patent Ductus Arteriosus, PS: Pulmonary Stenosis, RV: Right Ventricle, TOF: Tetralogy of Fallot, VSD: Ventricular Septal Defect.

Among 106 EA patients studied between 2001 and 2003 at the Mayo Clinic, there were 76 severe cases, 19 (18%) presented with LVNC, and 22 (21%) with WPW syndrome [78]. While no correlation existed between anatomical EA severity with the degree of LVNC, the QRS axis mean, which depicts ventricular depolarization on an electrocardiogram (ECG), showed a marginally significant difference (12° mean in EA/LVNC patients when compared with the 36° of all patients) [78].

In a study of 12 survivors among 16 neonates with EA who underwent right ventricular exclusion surgery, shortening fractions were found to be significantly lower prior to follow-up Glenn and Fontan procedures [7]. In our review, arrhythmia is a common clinical sign of EA/LVNC that appears second only to exertional dyspnea.

3.4. ECHO Is Preferred for EA Diagnosis While Cardiac MRI Is Preferred for LVNC

All imaging modalities used for diagnosis and EA/LVNC characterization are noted in Table 2. Imaging via ECHO is preferred for both CHDs, as it is noninvasive and cost-friendly [92]. However, LVNC diagnosis using ECHO is controversial because of a perceived arbitrary interpretation in imaging [15]. Cardiac MRI is thus recommended for diagnosing LVNC [45,56]. For quantitative assessment of volumetric and functional fractions, a cardiac CT may be used. Additional methods that offer further insight but less specificity and sensitivity for EA and LVNC include ECG, pulse oximetry, and exercise tests. In our review, we found most studies presented imaging and figures from 2-D ECHO, color flow doppler ECHO, cardiac MRI, and an ECG (Table 2).

3.5. EA/LVNC Treatment Is Non-Specific and Heterogenous

EA/LVNC treatment (Table 2) is nonspecific and ranges from symptomatic management to surgical valve repair [93,94]. Commonly, treatment included medication for heart failure or anticoagulation. Mechanical ventilation was common for earlier or severe presentation. Surgical repair often comprised arterioplasty, annuloplasty, closure of septal defects, and transplantation. Arrhythmia treatment consisted of medication or devices such as an implantable cardioverter-defibrillator (ICD) or left ventricular assist device (LVAD). In one CHD and/or LVNC-associated ICD study, EA patients suffered the highest complication rate during device implant, replacement, or lead-related procedures [86]. Most case reports, however, neglected details of the treatment initiated or discussed case prognosis.

3.6. Most Clinical Trials of EA or LVNC Seek to Assess CHD Genetic Etiology

All clinical trial information available for EA or LVNC in the US and EU are described in Table S1 (Supplementary Materials). Most were prospective observational studies and mainly sought to stratify risk and diagnostic criteria for each CHD. Some also studied the genetic underpinnings of EA or LVNC. One active trial (NCT02914171) investigating a nonrandomized therapeutic intervention of autologous bone marrow-derived mononuclear cells to treat 10 EA patients is projected to complete soon and may offer insight into EA therapies outside of the common medication or cardiac surgical procedures. Two other trials (NCT02432092 and NCT04265040) actively recruiting patients with cardiomyopathy (including LVNC) are studying molecular genetics and all-cause mortality outcomes.

3.7. Sarcomeric Variants Are Implicated in the Genetic Etiology of EA/LVNC

The genetic etiology and pathogenesis of EA/LVNC are largely unknown. Studies identified clinically significant variants at a variety of loci in EA and LVNC (*MYH7* [68,69,76], *NKX2.5* [25,95], *GATA4* [96], *TAZ* [97], *TBX20* [98], and others [74,99–102]), but few have been reported in EA/LVNC (Table 4). While independent EA- [103] or LVNC-associated [83] genetic variants have been discussed in the literature, Table 4 summarizes all genetic variant associations for EA/LVNC to date.

Many of these variants are in sarcomeric or cytoskeletal proteins and have broad implications in cardiac formation, function, and conduction [71,103]. The *MYH7* locus, which

encodes the β -myosin heavy chain (β MyHC), is significantly implicated in EA, LVNC, and EA/LVNC [68–70,75–77,83,87,88,104]. EA/LVNC-specific studies are detailed below.

Table 4. Summary of published work describing combined EA/LVNC-associated variants.

Year	Publication	Sample Size	Occurrence	Findings
2007	Budde et al. [69]	24	Familial	<i>MYH7</i> (p.R281T)
2010	Hoedemaekers et al. [104]	58	Sporadic	<i>MYH7</i> (p.L301Q)
2011	Postma et al. [76]	141	Sporadic	<i>MYH7</i> ; 7 mutations (5 novel)
2014	Hirono et al. [70]	3	Familial	<i>MYH7</i> (p.M362R)
2016	Kelle et al. [50]	1	Sporadic	<i>TPM1</i> (p.D159)
2018	Nijak et al. [71]	5	Familial	<i>TPM1</i> (p.L131V)
2019	Carlston et al. [41]	1	Sporadic	<i>NONO</i> (p.N52Sfs)
2020	Hirono et al. [83]	53	Sporadic	30 genetic variants in <i>MYH7</i> , <i>TPM1</i> , <i>ACTC1</i> , <i>ANK2</i> , <i>COL4A1</i> , <i>DAAM1</i> , <i>DSG2</i> , <i>DSP</i> , <i>FGF16</i> , <i>FGFR2</i> , <i>HCN4</i> , <i>JUP</i> , <i>MYBPC3</i> , <i>MYH6</i> , <i>MYL2</i> , <i>PKP2</i> , <i>PRDM16</i> , <i>RYR2</i> , and <i>TAZ</i>
2020	Samudrala et al. [72]	17	Familial	<i>KLHL26</i> (p.R237C)
2021	Mehdi et al. [56]	1	Sporadic	Gain chromosome band 15q11.2 and 1q44
2022	Coetzer et al. [42]	1	Sporadic	<i>NONO</i> (p.M389_T400del)
2022	Tu et al. [75]	6	Familial	<i>MYH7</i> splicing variant

Following a genome-wide linkage analysis of a German familial case of LVNC with 11 affected members among 25, Budde et al. discovered the *MHY7* (p.R281T) variant. Of the 11 LVNC-affected members, 4 also presented with EA. This *MYH7* variant is suspected to prevent salt-bridge formation, destabilizing the myosin head and resulting in altered contractility [69].

Postma et al. examined the *MYH7* locus of 141 unrelated EA patients via mutation screening and detected seven distinct heterozygous mutations, of which five were novel. In total, 6 patients among the 8 who possessed these *MYH7* variants and none of the remaining 133 subjects presented with LVNC [76].

After reviewing this latter publication and a few other EA/LVNC genetic studies, Vermeer et al. concluded the association of at least nine distinct *MYH7* mutations with EA/LVNC [87]. Five of these are located within the head region of β MyHC, while the other four reside in the rod domain, which mediates filament assembly and the sliding motion of contraction. Vermeer et al. further describe a subtype of EA/LVNC resulting from *MYH7* mutations or possibly other sarcomeric variants that follow a pattern of autosomal dominant inheritance and variable penetrance [87]. This conclusion has since been supported by isolated case reports [70] and variants found in other sarcomeric proteins such as thin filament α -tropomyosin, *TPM1* (p.D159N and p.L131V) [50,71].

Most recently, in a familial case of LVNC where only the proband presented with EA/LVNC, exome sequencing of the proband and parents and subsequent Sanger sequencing of other family members identified a maternally inherited heterozygous splicing variant in *MYH7* [75].

Non-sarcomeric proteins linked to EA/LVNC include *SCN5A*, a sodium channel, *NONO*, a nonoctamer-containing POU-domain DNA-binding protein, and *KLHL26*, a kelch-like protein (Table 4).

Neu et al. described one EA/LVNC-affected member among four related children who presented with early cardiac arrhythmia; all four were heterozygous for the *SCN5A* (p.I230T) variant [74].

The NONO protein plays crucial roles in transcriptional regulation and RNA splicing. One frameshift and one deletion variant of NONO were discovered in EA/LVNC patients who also presented with a noncardiac syndrome [41,42].

In a separate case, one EA/LVNC patient tested negative for a panel of 17 genes, including *TPM1* and *MYH7*, but showed chromosomal gain bands in 15q11.2 and 1q44 [56].

Recently, our group identified 10 EA/LVNC-affected members out of 17 in a familial study with an autosomal dominant pattern of inheritance [72]. From exome sequencing and subsequent Sanger sequencing, we discovered that the *KLHL26* (p.R237C) variant segregated with disease. The KLHL family of proteins is known to have a role in actin binding, cytoskeletal rearrangement, and ubiquitin-mediated protein degradation.

The abundance of loci implicated in either EA or LVNC alone and combined EA/LVNC underscores the complexity and genetic uncertainty for these CHDs.

4. Discussion

The characterization of EA as a distinct right heart defect persisted for nearly 150 years since its discovery in 1866 by Wilhelm Ebstein. Similarly, LVNC is mainly defined as a condition of the left heart. However, in the last 10–15 years, there has been an accumulation of evidence for more than a coincidental occurrence of EA and LVNC. Because of the prevalence of right-hearted lesions in LVNC and left-hearted lesions in EA seen in this review and others [78,105], a thorough review of the whole heart is recommended during workup of EA and/or LVNC.

4.1. Shared Developmental Pathways of EA and LVNC

The human heart begins to develop during the third week of human gestation [106]. The heart tube initially consists of an outer myocardial layer surrounded by the epicardium and an inner endocardial layer separated by cardiac jelly (extracellular matrix) (Figure 3A) [107]. Starting in week 4, a subset of overlying endocardial cells undergo endocardial-to-mesenchymal transformation (EndEMT) and invade the cardiac jelly overlying the atrioventricular canal, and the outflow tract swells to form the cardiac cushions (Figure 3B) [108,109]. In weeks 6–8, the cushions undergo extensive remodeling to form the mitral and tricuspid valves in the atrioventricular canal [110,111] (Figure 3C,D) and semi-lunar valves in the developing aorta and pulmonary trunk regions [109,111].

Cardiac fibroblasts within the developing myocardium are largely derived following epicardial-to-mesenchymal transition (EMT) of the epicardium and are known, in parallel, to provide signals for development of the atrioventricular valve as well as the compact myocardium [26,112]. The processes of EndEMT and EMT are governed by multiple pathways including the TGF- β /Smad, Wnt/ β -catenin, Notch, and Smad-independent TGF- β pathways [113,114]. Regulatory factors such as bone morphogenetic proteins (BMPs) and fibroblast growth factors (FGFs) prevent valve hyper- or hypoplasia [107,115].

The tricuspid valve is comprised of anterior, septal, and posterior leaflets; delamination of the septal leaflet from the muscular ventricular septum [116,117] likely occurs via anoikis (integrin-mediated apoptosis) [115]. Failure of delamination is thought to result in EA, which is characterized by an apically displaced tricuspid valve with a varied degree of freedom in movement (Figure 3E).

As these events unfold, myocardial compaction of the trabeculae, which began forming in week 4, occurs in week 8 [110,118]. As the myocardium expands via cardiomyocyte proliferation, endocardial cells invaginate, forming sheet-like protrusions (Figure 3F,G) comprising the trabeculae (light purple in Figure 3G), which is theorized to facilitate exchange of oxygen and nutrients [14]. By mid-gestation, the trabeculae become compacted (Figure 3H) [8,119]. Important factors in early trabeculation are a result of signaling via

endocardial and myocardial Notch, myocardial BMPs, epicardial FGFs, and both canonical and noncanonical Wnts [14].

Two LVNC investigations offered direct insight into genetic etiology. One described autosomal dominant LVNC associated with a *MIB1* (mindbomb homolog 1) variant [101]. *MIB1* encodes an E3-ubiquitin ligase that promotes endocytosis of two NOTCH ligands. The second study modeled LVNC associated with a variant in transcription factor *TBX20* using induced pluripotent stem cells (iPSCs); they found decreased cellular proliferation in disease lines and showed phenotype reversal following variant correction [98].




Figure 3. Development of the tricuspid valve and compaction of the ventricular myocardium: Cardiogenesis is initiated as (A) heart tube consisting of an outer epicardium that surrounds a layer of myocardial cells, with inner cardiac jelly and an endocardial cell layer; (B) following, the heart tube undergoes rightward looping, and endocardial cells overlying the primitive valve regions undergo endothelial-to-mesenchymal transformation and invade the cardiac jelly to form swellings termed endocardial cushions; (C) these cushions then remodel and elongate into leaflets/cusps and supporting structures. In the atrioventricular position (mitral, tricuspid), the leaflets separate from the myocardial wall via delamination and (D) become freely movable; (E) an example of an apically displaced valve leaflet with incomplete separation from the myocardial wall typically seen in EA; (F) cardiomyocyte proliferation and endocardial cell invagination form the sheet-like protrusions of the (G) trabecular layer seen in light purple; (H) later, this trabecular layer becomes compacted. Parallel signaling from TGF- β /Smad, Wnt/ β -catenin, noncanonical Wnt, Notch, and Smad-independent TGF- β , BMP, and FGF pathways enable the development of both the tricuspid valve and compaction of the ventricular myocardium. Time indicated corresponds with human development.

4.2. A Common Etiology for EA and LVNC?

While EA/LVNC pathophysiology remains unknown, when we reflect on the considerable overlap of developmental pathways in the formation of the tricuspid valve and compacted myocardium and the genetic variants associated with both CHDs, it is unsurprising that EA commonly manifests with LVNC.

Investigating sarcomeric and cytoskeletal variants may provide evidence for arrhythmia stemming from a cellular or tissue-level contractile impediment rather than a global consequence of anatomical alterations. Aberrant contractile ability may further hinder apical constriction, depolarization, and subsequent delamination of cells during EndEMT or EMT processes [120].

5. Conclusions

Understanding the molecular and cellular events that unfold during the progression of disease-specific variants will provide broader insight into typical vs. aberrant cardiac development. Although clinical approaches have improved, there remains an unmet need for effective EA/LVNC treatment. Nonspecific pharmaceutical and mechanical interventions will likely remain the most common frontline treatment options for EA/LVNC until a deeper understanding of the genetic and molecular underpinnings of both CHDs becomes available.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/jcdd9040115/s1>, Table S1: Summary of Clinical Trials on EA, LVNC, or EA/LVNC.

Author Contributions: Conceptualization, S.K.T.; writing—original draft preparation, S.K.T.; writing—review and editing, S.K.T., M.A.F., J.L., J.W.L., M.E.M. and A.T.-M.; supervision, M.E.M. and A.T.-M. All authors have read and agreed to the published version of the manuscript.

Funding: S.K.T. is a member of the Medical Scientist Training Program at MCW, which is partially supported by a training grant from the National Institute of General Medical Sciences, T32-GM080202. Additional funding came from the Children’s Wisconsin Research Institute and the MCW Department of Surgery.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Attenhofer Jost, C.H.; Connolly, H.M.; Dearani, J.A.; Edwards, W.D.; Danielson, G.K. Ebstein’s anomaly. *Circulation* **2007**, *115*, 277–285. [[CrossRef](#)] [[PubMed](#)]
- Correa-Villasenor, A.; Ferencz, C.; Neill, C.A.; Wilson, P.D.; Boughman, J.A. Ebstein’s malformation of the tricuspid valve: Genetic and environmental factors. The Baltimore-Washington Infant Study Group. *Teratology* **1994**, *50*, 137–147. [[CrossRef](#)] [[PubMed](#)]
- Lupo, P.J.; Langlois, P.H.; Mitchell, L.E. Epidemiology of Ebstein anomaly: Prevalence and patterns in Texas, 1999–2005. *Am. J. Med. Genet. A* **2011**, *155*, 1007–1014. [[CrossRef](#)]
- Fahed, A.C.; Gelb, B.D.; Seidman, J.G.; Seidman, C.E. Genetics of congenital heart disease: The glass half empty. *Circ. Res.* **2013**, *112*, 707–720. [[CrossRef](#)] [[PubMed](#)]
- Ho, S.Y.; Goltz, D.; McCarthy, K.; Cook, A.C.; Connell, M.G.; Smith, A.; Anderson, R.H. The atrioventricular junctions in Ebstein malformation. *Heart* **2000**, *83*, 444–449. [[CrossRef](#)]
- Chauvaud, S. Ebstein’s malformation. surgical treatment and results. *Thorac. Cardiovasc. Surg.* **2000**, *48*, 220–223. [[CrossRef](#)] [[PubMed](#)]
- Reemtsen, B.L.; Polimenakos, A.C.; Fagan, B.T.; Wells, W.J.; Starnes, V.A. Fate of the right ventricle after fenestrated right ventricular exclusion for severe neonatal Ebstein anomaly. *J. Thorac. Cardiovasc. Surg.* **2007**, *134*, 1406–1410. [[CrossRef](#)] [[PubMed](#)]
- Freedom, R.M.; Yoo, S.J.; Perrin, D.; Taylor, G.; Petersen, S.; Anderson, R.H. The morphological spectrum of ventricular noncompaction. *Cardiol. Young* **2005**, *15*, 345–364. [[CrossRef](#)] [[PubMed](#)]
- Voges, I.; Al-Mallah, M.H.; Scognamiglio, G.; Di Salvo, G. Right Heart-Pulmonary Circulation Unit in Congenital Heart Diseases. *Heart Fail. Clin.* **2018**, *14*, 283–295. [[CrossRef](#)]
- Celermajer, D.S.; Bull, C.; Till, J.A.; Cullen, S.; Vassilikos, V.P.; Sullivan, I.D.; Allan, L.; Nihoyannopoulos, P.; Somerville, J.; Deanfield, J.E. Ebstein’s anomaly: Presentation and outcome from fetus to adult. *J. Am. Coll. Cardiol.* **1994**, *23*, 170–176. [[CrossRef](#)]
- Cabrera, R.; Miranda-Fernandez, M.C.; Huertas-Quinones, V.M.; Carreno, M.; Pineda, I.; Restrepo, C.M.; Silva, C.T.; Quero, R.; Cano, J.D.; Manrique, D.C.; et al. Identification of clinically relevant phenotypes in patients with Ebstein anomaly. *Clin. Cardiol.* **2018**, *41*, 343–348. [[CrossRef](#)] [[PubMed](#)]
- Carpentier, A.; Chauvaud, S.; Mace, L.; Relland, J.; Mihaileanu, S.; Marino, J.P.; Abry, B.; Guibourt, P. A new reconstructive operation for Ebstein’s anomaly of the tricuspid valve. *J. Thorac. Cardiovasc. Surg.* **1988**, *96*, 92–101. [[CrossRef](#)]
- Dearani, J.A.; Danielson, G.K. Congenital Heart Surgery Nomenclature and Database Project: Ebstein’s anomaly and tricuspid valve disease. *Ann. Thorac. Surg.* **2000**, *69*, S106–S117. [[CrossRef](#)]

14. Zhang, W.; Chen, H.; Qu, X.; Chang, C.P.; Shou, W. Molecular mechanism of ventricular trabeculation/compaction and the pathogenesis of the left ventricular noncompaction cardiomyopathy (LVNC). *Am. J. Med. Genet. C Semin. Med. Genet.* **2013**, *163*, 144–156. [[CrossRef](#)] [[PubMed](#)]
15. Towbin, J.A.; Lorts, A.; Jefferies, J.L. Left ventricular non-compaction cardiomyopathy. *Lancet* **2015**, *386*, 813–825. [[CrossRef](#)]
16. Tarando, F.; Coisne, D.; Galli, E.; Rousseau, C.; Viera, F.; Bosseau, C.; Habib, G.; Lederlin, M.; Schnell, F.; Donal, E. Left ventricular non-compaction and idiopathic dilated cardiomyopathy: The significant diagnostic value of longitudinal strain. *Int. J. Cardiovasc. Imaging* **2017**, *33*, 83–95. [[CrossRef](#)]
17. Tsai, S.F.; Ebenroth, E.S.; Hurwitz, R.A.; Cordes, T.M.; Schamberger, M.S.; Batra, A.S. Is left ventricular noncompaction in children truly an isolated lesion? *Pediatr. Cardiol.* **2009**, *30*, 597–602. [[CrossRef](#)]
18. El-Menyar, A.A.; Gendi, S.M.; Numan, M.T. Noncompaction cardiomyopathy in the State of Qatar. *Saudi Med. J.* **2007**, *28*, 429–434.
19. Chin, T.K.; Perloff, J.K.; Williams, R.G.; Jue, K.; Mohrmann, R. Isolated noncompaction of left ventricular myocardium. A study of eight cases. *Circulation* **1990**, *82*, 507–513. [[CrossRef](#)]
20. Jenni, R.; Oechslin, E.; Schneider, J.; Attenhofer Jost, C.; Kaufmann, P.A. Echocardiographic and pathoanatomical characteristics of isolated left ventricular non-compaction: A step towards classification as a distinct cardiomyopathy. *Heart* **2001**, *86*, 666–671. [[CrossRef](#)]
21. Petersen, S.E.; Selvanayagam, J.B.; Wiesmann, F.; Robson, M.D.; Francis, J.M.; Anderson, R.H.; Watkins, H.; Neubauer, S. Left ventricular non-compaction: Insights from cardiovascular magnetic resonance imaging. *J. Am. Coll. Cardiol.* **2005**, *46*, 101–105. [[CrossRef](#)] [[PubMed](#)]
22. Attenhofer Jost, C.H.; Connolly, H.M.; Warnes, C.A.; O’Leary, P.; Tajik, A.J.; Pellikka, P.A.; Seward, J.B. Noncompacted myocardium in Ebstein’s anomaly: Initial description in three patients. *J. Am. Soc. Echocardiogr.* **2004**, *17*, 677–680. [[CrossRef](#)] [[PubMed](#)]
23. Page, M.J.; McKenzie, J.E.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ* **2021**, *372*, n71. [[CrossRef](#)]
24. Page, M.J.; Moher, D.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. PRISMA 2020 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews. *BMJ* **2021**, *372*, n160. [[CrossRef](#)] [[PubMed](#)]
25. Ashraf, H.; Pradhan, L.; Chang, E.I.; Terada, R.; Ryan, N.J.; Briggs, L.E.; Chowdhury, R.; Zarate, M.A.; Sugi, Y.; Nam, H.J.; et al. A mouse model of human congenital heart disease: High incidence of diverse cardiac anomalies and ventricular noncompaction produced by heterozygous Nkx2-5 homeodomain missense mutation. *Circ. Cardiovasc. Genet.* **2014**, *7*, 423–433. [[CrossRef](#)] [[PubMed](#)]
26. Gittenberger-de Groot, A.C.; Winter, E.M.; Poelmann, R.E. Epicardium-derived cells (EPDCs) in development, cardiac disease and repair of ischemia. *J. Cell Mol. Med.* **2010**, *14*, 1056–1060. [[CrossRef](#)]
27. Abu-Issa, R.; Waldo, K.; Kirby, M.L. Heart fields: One, two or more? *Dev. Biol.* **2004**, *272*, 281–285. [[CrossRef](#)] [[PubMed](#)]
28. Basson, C.T.; Bachinsky, D.R.; Lin, R.C.; Levi, T.; Elkins, J.A.; Soultz, J.; Grayzel, D.; Kroumpouzou, E.; Traill, T.A.; Leblanc-Straceski, J.; et al. Mutations in human TBX5 [corrected] cause limb and cardiac malformation in Holt-Oram syndrome. *Nat. Genet.* **1997**, *15*, 30–35. [[CrossRef](#)]
29. Al-Wakeel-Marquard, N.; Seidel, F.; Kuhnisch, J.; Kuehne, T.; Berger, F.; Messroghli, D.R.; Klaassen, S. Midwall Fibrosis and Cardiac Mechanics: Rigid Body Rotation Is a Novel Marker of Disease Severity in Pediatric Primary Dilated Cardiomyopathy. *Front. Cardiovasc. Med.* **2021**, *8*, 810005. [[CrossRef](#)]
30. Al-Wakeel-Marquard, N.; Degener, F.; Herbst, C.; Kuhnisch, J.; Dartsch, J.; Schmitt, B.; Kuehne, T.; Messroghli, D.; Berger, F.; Klaassen, S. RIKADA Study Reveals Risk Factors in Pediatric Primary Cardiomyopathy. *J. Am. Heart Assoc.* **2019**, *8*, e012531. [[CrossRef](#)]
31. Renilla, A.; Santamarta, E.; Corros, C.; Martin, M.; Barreiro, M.; de la Hera, J. Isolated left ventricular non-compaction associated with Ebstein’s anomaly. Multimodality non-invasive imaging for the assessment of congenital heart disease. *Arch. Cardiol. Mex.* **2013**, *83*, 100–103. [[CrossRef](#)] [[PubMed](#)]
32. Dima-Cozma, C.; Cojocar, D.C.; Chiriac, S.; Negru, R.; Mitu, F. Ebstein’s anomaly. *Rev. Med. Chir. Soc. Med. Nat. Iasi* **2013**, *117*, 674–679.
33. O’Leary, P.W.; Qureshi, M.Y.; Cetta, F.; Nelson, T.J.; Holst, K.A.; Dearani, J.A.; Wanek Program Clinical Pipeline, G. Cone Reconstruction for Ebstein Anomaly: Ventricular Remodeling and Preliminary Impact of Stem Cell Therapy. *Mayo Clin. Proc.* **2021**, *96*, 3053–3061. [[CrossRef](#)] [[PubMed](#)]
34. Tulino, V.; Cacace, C.; Tulino, D.; Imbalzano, E.; Dattilo, G. Clinical variants in Ebstein’s anomaly. *Int. J. Cardiol.* **2013**, *168*, 4969–4970. [[CrossRef](#)] [[PubMed](#)]
35. Andresen, H.; Kaag, N.; Potratz, J. Non-compaction of ventricular myocardium and contrast-enhanced echocardiography. *Z Kardiol.* **2005**, *94*, 483–485. [[CrossRef](#)] [[PubMed](#)]
36. Arslan, S.; Gurlertop, H.Y.; Gundogdu, F.; Senocak, H. Left ventricular noncompaction and mid-caviter narrowing associated with Ebstein’s anomaly: Three-dimensional transthoracic echocardiographic image. *Int. J. Cardiol.* **2007**, *115*, e52–e55. [[CrossRef](#)] [[PubMed](#)]
37. Aydin, C.; Engin, M. Ebstein’s anomaly associated with biventricular noncompaction. *J. Echocardiogr.* **2021**. [[CrossRef](#)] [[PubMed](#)]

38. Bagur, R.H.; Lederlin, M.; Montaudon, M.; Latrabe, V.; Corneloup, O.; Iriart, X.; Laurent, F. Images in cardiovascular medicine. Ebstein anomaly associated with left ventricular noncompaction. *Circulation* **2008**, *118*, e662–e664. [[CrossRef](#)]
39. Baysan, O.; Yokusoglu, M.; Bugan, B.; Demirkol, S. E-page original images. A case of Ebstein anomaly and biventricular noncompaction. *Anadolu Kardiyol. Derg.* **2010**, *10*, E26. [[CrossRef](#)]
40. Betrian Blasco, P.; Gallardo Agromayor, E. Ebstein's anomaly and left ventricular noncompaction association. *Int. J. Cardiol.* **2007**, *119*, 264–265. [[CrossRef](#)]
41. Carlston, C.M.; Bleyl, S.B.; Andrews, A.; Meyers, L.; Brown, S.; Bayrak-Toydemir, P.; Bale, J.F.; Botto, L.D. Expanding the genetic and clinical spectrum of the NONO-associated X-linked intellectual disability syndrome. *Am. J. Med. Genet. A* **2019**, *179*, 792–796. [[CrossRef](#)]
42. Coetzer, K.C.; Moosa, S. Novel hemizygous loss-of-function variant in NONO identified in a South African boy. *Am. J. Med. Genet. A* **2022**, *188*, 373–376. [[CrossRef](#)]
43. Das, B.B.; Chrisant, M.; Lavandosky, G.; Zakrzewski, M.; Winchester, R.; Turner, I.; Bibevski, S.; Scholl, F. An Adolescent with Left Ventricular Noncompaction and Ebstein Anomaly Presenting with Advanced Heart Failure: Discharge from Hospital with a Biventricular Assist Device. *J. Pediatr.* **2018**, *202*, 304–310. [[CrossRef](#)]
44. Dogan, A.; Aksoy, H. Coexistence of congenital heart anomalies with noncompaction of the ventricular myocardium. Two case reports. *Herz* **2012**, *37*, 699–701. [[CrossRef](#)]
45. Evans, E.; McDonald, M.; Adamson, D.L.; Khan, J.N. Cardiomyopathy in the peripartum period due to left ventricular non-compaction and association with Ebstein's anomaly: A case report. *Eur Heart J. Case Rep.* **2021**, *5*, ytab088. [[CrossRef](#)]
46. Fazio, G.; Visconti, C.; D'Angelo, L.; Grassedonio, E.; Lo Re, G.; D'Amico, T.; Sutura, L.; Novo, G.; Ferrara, F.; Midiri, M.; et al. Diagnosis and definition of biventricular non-compaction associated to Ebstein's anomaly. *Int. J. Cardiol.* **2011**, *150*, e20–e24. [[CrossRef](#)]
47. Geerdink, L.M.; Ter Heide, H.; Kapusta, L. Ebstein's anomaly and aortic arch lesions: When right meets left. *Int. J. Cardiol.* **2016**, *223*, 568–570. [[CrossRef](#)]
48. Ilercil, A.; Barack, J.; Malone, M.A.; Barold, S.S.; Herweg, B. Association of noncompaction of left ventricular myocardium with Ebstein's anomaly. *Echocardiography* **2006**, *23*, 432–433. [[CrossRef](#)]
49. Isilak, Z.; Tokatli, A.; Yalcin, M.; Uzun, M. Case images: A very rare case: Ebstein anomaly, biventricular noncompaction and pre-excitation in the same patient. *Turk Kardiyol. Dern. Ars.* **2012**, *40*, 557. [[CrossRef](#)]
50. Kelle, A.M.; Bentley, S.J.; Rohena, L.O.; Cabalka, A.K.; Olson, T.M. Ebstein anomaly, left ventricular non-compaction, and early onset heart failure associated with a de novo alpha-tropomyosin gene mutation. *Am. J. Med. Genet. A* **2016**, *170*, 2186–2190. [[CrossRef](#)]
51. Kharge, J.; Prasad, M.R.; Ramegowda, R.T. An unusual case of congenitally corrected transposition of the great arteries associated with noncompaction-like remodeling of the morphological right ventricle. *Echocardiography* **2011**, *28*, E212–E214. [[CrossRef](#)]
52. Kiernan, T.J.; Fahy, G. Multiple accessory pathways, dual AV nodal physiology, non-compacted myocardium and patent foramen ovale in a patient with Ebstein's anomaly: Report of a case. *Int. J. Cardiol.* **2007**, *114*, 412–413. [[CrossRef](#)]
53. Kumar, N.; Troianos, C.A.; Baisden, J.S. Left Ventricular Assist Device Insertion in a Patient With Biventricular Noncompaction Cardiomyopathy, Ebstein Anomaly, and a Left Atrial Mass: A Case Report. *A & A Case Rep.* **2016**, *7*, 251–255. [[CrossRef](#)]
54. Malagoli, A.; Rossi, L.; Mastrojanni, C.; Villani, G.Q. A perfect storm: Wolf Parkinson White syndrome, Ebstein's anomaly, biventricular non-compaction, and bicuspid aortic valve. *Eur. Heart J. Cardiovasc. Imaging* **2014**, *15*, 827. [[CrossRef](#)]
55. McGee, M.; Warner, L.; Collins, N. Ebstein's Anomaly, Left Ventricular Noncompaction, and Sudden Cardiac Death. *Case Rep. Cardiol.* **2015**, *2015*, 854236. [[CrossRef](#)]
56. Mehdi, M.; Bhatia, S.; Patel, M.; Aly, A. Ebstein's Anomaly, Left Ventricular Noncompaction and Gerbode-Like Defect Triad (Fetal Diagnosis and Neonatal Course). *Case Rep. Pediatr.* **2021**, *2021*, 9969588. [[CrossRef](#)]
57. Nimeri, N.A.; Abou Nahia, F.F.; Ibrahim, A.S.; Khella, A.Y. The first reported case of non-compacted cardiomyopathy in a preterm infant with Ebstein's anomaly. *BMJ Case Rep.* **2012**, *2012*, bcr0220125861. [[CrossRef](#)]
58. Patra, S.; Singla, V.; Kharge, J.; Ravindranath, K.S.; Manjunath, C.N. A patient of Ebstein's anomaly associated with biventricular noncompaction presented with Wolf Parkinson White syndrome—A rare presentation. *J. Cardiovasc. Dis. Res.* **2012**, *3*, 323–325. [[CrossRef](#)]
59. Saltik, L.; Eroglu, E.; Bayrak, F.; Degertekin, M. Left ventricular noncompaction associated with Ebstein's anomaly. *Turk Kardiyol. Dern. Ars.* **2009**, *37*, 269–272.
60. Scott, D.A.; Hernandez-Garcia, A.; Azamian, M.S.; Jordan, V.K.; Kim, B.J.; Starkovich, M.; Zhang, J.; Wong, L.J.; Darilek, S.A.; Breman, A.M.; et al. Congenital heart defects and left ventricular non-compaction in males with loss-of-function variants in NONO. *J. Med. Genet.* **2017**, *54*, 47–53. [[CrossRef](#)]
61. Song, Z.Z. A combination of right ventricular hypertrabeculation/noncompaction and Ebstein's anomaly. *Int. J. Cardiol.* **2010**, *143*, e30–e33. [[CrossRef](#)] [[PubMed](#)]
62. Stollberger, C.; Kopsa, W.; Finsterer, J. Non-compaction of the right atrium and left ventricle in Ebstein's malformation. *J. Heart Valve Dis.* **2006**, *15*, 719–720. [[PubMed](#)]
63. Thakur, V.; Jaeggi, E.T.; Nield, L.E. A unique foetal case of left ventricular non-compaction associated with arrhythmia, structural cardiac anomalies, and agenesis of the ductus venosus. *Cardiol. Young* **2016**, *26*, 368–370. [[CrossRef](#)] [[PubMed](#)]

64. Thani, K.B.; Khadivi, B.; Kahn, A.M.; Cotter, B.; Blanchard, D. Ebstein's anomaly with left ventricular noncompaction and bicuspid aortic valve. *J. Am. Coll. Cardiol.* **2010**, *56*, 899. [[CrossRef](#)] [[PubMed](#)]
65. Tourmousoglou, C.; Bogossian, H.; Ninios, V.; Ninios, E. A rare case of Ebstein's anomaly with left ventricular noncompaction. *Asian Cardiovasc. Thorac. Ann.* **2019**, *27*, 208–209. [[CrossRef](#)] [[PubMed](#)]
66. Wijesekera, V.; Yadav, R. Combined left ventricular non-compaction and Ebstein's anomaly. *Heart Asia* **2012**, *4*, 37. [[CrossRef](#)] [[PubMed](#)]
67. Zarrouk Mahjoub, S.; Mehri, S.; Ourda, F.; Boussaada, R.; Mechmeche, R.; Arab, S.B.; Finsterer, J. Transition m.3308T>C in the ND1 gene is associated with left ventricular hypertrabeculation/noncompaction. *Cardiology* **2011**, *118*, 153–158. [[CrossRef](#)]
68. Bettinelli, A.L.; Mulder, T.J.; Funke, B.H.; Lafferty, K.A.; Longo, S.A.; Niyazov, D.M. Familial ebstein anomaly, left ventricular hypertrabeculation, and ventricular septal defect associated with a MYH7 mutation. *Am. J. Med. Genet. A* **2013**, *161*, 3187–3190. [[CrossRef](#)] [[PubMed](#)]
69. Budde, B.S.; Binner, P.; Waldmuller, S.; Hohne, W.; Blankenfeldt, W.; Hassfeld, S.; Bromsen, J.; Dermintzoglou, A.; Wieczorek, M.; May, E.; et al. Noncompaction of the ventricular myocardium is associated with a de novo mutation in the beta-myosin heavy chain gene. *PLoS ONE* **2007**, *2*, e1362. [[CrossRef](#)]
70. Hirono, K.; Hata, Y.; Ibuki, K.; Yoshimura, N. Familial Ebstein's anomaly, left ventricular noncompaction, and ventricular septal defect associated with an MYH7 mutation. *J. Thorac. Cardiovasc. Surg.* **2014**, *148*, e223–e226. [[CrossRef](#)]
71. Nijak, A.; Alaerts, M.; Kuiperi, C.; Corveleyn, A.; Suys, B.; Paelinck, B.; Saenen, J.; Van Craenenbroeck, E.; Van Laer, L.; Loeys, B.; et al. Left ventricular non-compaction with Ebstein anomaly attributed to a TPM1 mutation. *Eur. J. Med. Genet.* **2018**, *61*, 8–10. [[CrossRef](#)] [[PubMed](#)]
72. Samudrala, S.S.K.; North, L.M.; Stamm, K.D.; Earing, M.G.; Frommelt, M.A.; Willes, R.; Tripathi, S.; Dsouza, N.R.; Zimmermann, M.T.; Mahnke, D.K.; et al. Novel KLHL26 variant associated with a familial case of Ebstein's anomaly and left ventricular noncompaction. *Mol. Genet. Genom. Med.* **2020**, *8*, e1152. [[CrossRef](#)]
73. Sinkovec, M.; Kozelj, M.; Podnar, T. Familial biventricular myocardial noncompaction associated with Ebstein's malformation. *Int. J. Cardiol.* **2005**, *102*, 297–302. [[CrossRef](#)] [[PubMed](#)]
74. Neu, A.; Eiselt, M.; Paul, M.; Sauter, K.; Stallmeyer, B.; Isbrandt, D.; Schulze-Bahr, E. A homozygous SCN5A mutation in a severe, recessive type of cardiac conduction disease. *Hum. Mutat.* **2010**, *31*, E1609–E1621. [[CrossRef](#)] [[PubMed](#)]
75. Tu, P.; Sun, H.; Zhang, X.; Ran, Q.; He, Y.; Ran, S. Diverse cardiac phenotypes among different carriers of the same MYH7 splicing variant allele (c.732+1G>A) from a family. *BMC Med. Genom.* **2022**, *15*, 36. [[CrossRef](#)] [[PubMed](#)]
76. Postma, A.V.; van Engelen, K.; van de Meerakker, J.; Rahman, T.; Probst, S.; Baars, M.J.; Bauer, U.; Pickardt, T.; Sperling, S.R.; Berger, F.; et al. Mutations in the sarcomere gene MYH7 in Ebstein anomaly. *Circ. Cardiovasc. Genet.* **2011**, *4*, 43–50. [[CrossRef](#)] [[PubMed](#)]
77. van Engelen, K.; Postma, A.V.; van de Meerakker, J.B.; Roos-Hesselink, J.W.; Helderma-van den Enden, A.T.; Vliegen, H.W.; Rahman, T.; Baars, M.J.; Sels, J.W.; Bauer, U.; et al. Ebstein's anomaly may be caused by mutations in the sarcomere protein gene MYH7. *Neth Heart J.* **2013**, *21*, 113–117. [[CrossRef](#)]
78. Attenhofer Jost, C.H.; Connolly, H.M.; O'Leary, P.W.; Warnes, C.A.; Tajik, A.J.; Seward, J.B. Left heart lesions in patients with Ebstein anomaly. *Mayo Clin. Proc.* **2005**, *80*, 361–368. [[CrossRef](#)]
79. Stahli, B.E.; Gebhard, C.; Biaggi, P.; Klaassen, S.; Valsangiacomo Buechel, E.; Attenhofer Jost, C.H.; Jenni, R.; Tanner, F.C.; Greutmann, M. Left ventricular non-compaction: Prevalence in congenital heart disease. *Int. J. Cardiol.* **2013**, *167*, 2477–2481. [[CrossRef](#)]
80. Kumor, M.; Lipczynska, M.; Biernacka, E.K.; Klisiewicz, A.; Wojcik, A.; Konka, M.; Kozuch, K.; Szymanski, P.; Hoffman, P. Cardiac arrest and ventricular arrhythmia in adults with Ebstein anomaly and left ventricular non-compaction. *J. Cardiol.* **2018**, *71*, 484–487. [[CrossRef](#)]
81. Marques, L.C.; Liguori, G.R.; Amarante Souza, A.C.; Aiello, V.D. Left Ventricular Noncompaction Is More Prevalent in Ventricular Septal Defect Than Other Congenital Heart Defects: A Morphological Study. *J. Cardiovasc Dev. Dis* **2020**, *7*, 39. [[CrossRef](#)] [[PubMed](#)]
82. Pignatelli, R.H.; Texter, K.M.; Denfield, S.W.; Grenier, M.A.; Altman, C.A.; Ayres, N.A.; Chandra-Bose Reddy, S. LV Noncompaction in Ebstein's anomaly in infants and outcomes. *JACC Cardiovasc. Imaging* **2014**, *7*, 207–209. [[CrossRef](#)]
83. Hirono, K.; Hata, Y.; Miyao, N.; Okabe, M.; Takarada, S.; Nakaoka, H.; Ibuki, K.; Ozawa, S.; Yoshimura, N.; Nishida, N.; et al. Left Ventricular Noncompaction and Congenital Heart Disease Increases the Risk of Congestive Heart Failure. *J. Clin. Med.* **2020**, *9*, 785. [[CrossRef](#)] [[PubMed](#)]
84. Yuan, S.M. Ebstein's Anomaly: Genetics, Clinical Manifestations, and Management. *Pediatr. Neonatol.* **2017**, *58*, 211–215. [[CrossRef](#)] [[PubMed](#)]
85. Barra, S.N.; Providencia, R.; Paiva, L.; Nascimento, J.; Marques, A.L. A review on advanced atrioventricular block in young or middle-aged adults. *Pacing Clin. Electrophysiol.* **2012**, *35*, 1395–1405. [[CrossRef](#)]
86. Gleva, M.J.; Wang, Y.; Curtis, J.P.; Berul, C.I.; Huddleston, C.B.; Poole, J.E. Complications Associated With Implantable Cardioverter Defibrillators in Adults With Congenital Heart Disease or Left Ventricular Noncompaction Cardiomyopathy (From the NCDR(R) Implantable Cardioverter-Defibrillator Registry). *Am. J. Cardiol.* **2017**, *120*, 1891–1898. [[CrossRef](#)]

87. Vermeer, A.M.; van Engelen, K.; Postma, A.V.; Baars, M.J.; Christiaans, I.; De Haij, S.; Klaassen, S.; Mulder, B.J.; Keavney, B. Ebstein anomaly associated with left ventricular noncompaction: An autosomal dominant condition that can be caused by mutations in MYH7. *Am. J. Med. Genet. C Semin. Med. Genet.* **2013**, *163*, 178–184. [[CrossRef](#)]
88. Kayvanpour, E.; Sedaghat-Hamedani, F.; Gi, W.T.; Tugrul, O.F.; Amr, A.; Haas, J.; Zhu, F.; Ehlermann, P.; Uhlmann, L.; Katus, H.A.; et al. Clinical and genetic insights into non-compaction: A meta-analysis and systematic review on 7598 individuals. *Clin. Res. Cardiol.* **2019**, *108*, 1297–1308. [[CrossRef](#)]
89. Monibi, A.A.; Neches, W.H.; Lenox, C.C.; Park, S.C.; Mathews, R.A.; Zuberbuhler, J.R. Left ventricular anomalies associated with Ebstein's malformation of the tricuspid valve. *Circulation* **1978**, *57*, 303–306. [[CrossRef](#)] [[PubMed](#)]
90. Daliendo, L.; Angelini, A.; Ho, S.Y.; Frescura, C.; Turrini, P.; Baratella, M.C.; Thiene, G.; Anderson, R.H. Angiographic and morphologic features of the left ventricle in Ebstein's malformation. *Am. J. Cardiol.* **1997**, *80*, 1051–1059. [[CrossRef](#)]
91. Chin, K.T.; Xu, H.T.; Ching, Y.P.; Jin, D.Y. Differential subcellular localization and activity of kelch repeat proteins KLHDC1 and KLHDC2. *Mol. Cell Biochem.* **2007**, *296*, 109–119. [[CrossRef](#)]
92. Qureshi, M.Y.; O'Leary, P.W.; Connolly, H.M. Cardiac imaging in Ebstein anomaly. *Trends Cardiovasc. Med.* **2018**, *28*, 403–409. [[CrossRef](#)] [[PubMed](#)]
93. Kim, M.S.; Lim, H.G.; Kim, W.H.; Lee, J.R.; Kim, Y.J. Long-Term Results after Surgical Treatment of Ebstein's Anomaly: A 30-year Experience. *Korean Circ. J.* **2016**, *46*, 706–713. [[CrossRef](#)] [[PubMed](#)]
94. Sainathan, S.; da Fonseca da Silva, L.; da Silva, J.P. Ebstein's anomaly: Contemporary management strategies. *J. Thorac. Dis.* **2020**, *12*, 1161–1173. [[CrossRef](#)] [[PubMed](#)]
95. Gioli-Pereira, L.; Pereira, A.C.; Mesquita, S.M.; Xavier-Neto, J.; Lopes, A.A.; Krieger, J.E. NKX2.5 mutations in patients with non-syndromic congenital heart disease. *Int. J. Cardiol.* **2010**, *138*, 261–265. [[CrossRef](#)]
96. Digilio, M.C.; Bernardini, L.; Lepri, F.; Giuffrida, M.G.; Guida, V.; Baban, A.; Versacci, P.; Capolino, R.; Torres, B.; De Luca, A.; et al. Ebstein anomaly: Genetic heterogeneity and association with microdeletions 1p36 and 8p23.1. *Am. J. Med. Genet. A* **2011**, *155*, 2196–2202. [[CrossRef](#)]
97. Bione, S.; D'Adamo, P.; Maestrini, E.; Gedeon, A.K.; Bolhuis, P.A.; Toniolo, D. A novel X-linked gene, G4.5, is responsible for Barth syndrome. *Nat. Genet.* **1996**, *12*, 385–389. [[CrossRef](#)]
98. Kodo, K.; Ong, S.G.; Jahanbani, F.; Termglinchan, V.; Hirono, K.; InanlooRahatloo, K.; Ebert, A.D.; Shukla, P.; Abilez, O.J.; Churko, J.M.; et al. iPSC-derived cardiomyocytes reveal abnormal TGF-beta signalling in left ventricular non-compaction cardiomyopathy. *Nat. Cell Biol.* **2016**, *18*, 1031–1042. [[CrossRef](#)] [[PubMed](#)]
99. van Waning, J.I.; Caliskan, K.; Hoedemaekers, Y.M.; van Spaendonck-Zwarts, K.Y.; Baas, A.F.; Boekholdt, S.M.; van Melle, J.P.; Teske, A.J.; Asselbergs, F.W.; Backx, A.; et al. Genetics, Clinical Features, and Long-Term Outcome of Noncompaction Cardiomyopathy. *J. Am. Coll. Cardiol.* **2018**, *71*, 711–722. [[CrossRef](#)]
100. Kenton, A.B.; Sanchez, X.; Coveler, K.J.; Makar, K.A.; Jimenez, S.; Ichida, F.; Murphy, R.T.; Elliott, P.M.; McKenna, W.; Bowles, N.E.; et al. Isolated left ventricular noncompaction is rarely caused by mutations in G4.5, alpha-dystrobrevin and FK Binding Protein-12. *Mol. Genet. Metab.* **2004**, *82*, 162–166. [[CrossRef](#)]
101. Luxan, G.; Casanova, J.C.; Martinez-Poveda, B.; Prados, B.; D'Amato, G.; MacGrogan, D.; Gonzalez-Rajal, A.; Dobarro, D.; Torroja, C.; Martinez, F.; et al. Mutations in the NOTCH pathway regulator MIB1 cause left ventricular noncompaction cardiomyopathy. *Nat. Med.* **2013**, *19*, 193–201. [[CrossRef](#)]
102. Mercer, C.L.; Andreoletti, G.; Carroll, A.; Salmon, A.P.; Temple, I.K.; Ennis, S. Familial Ebstein Anomaly: Whole Exome Sequencing Identifies Novel Phenotype Associated With FLNA. *Circ. Cardiovasc. Genet.* **2017**, *10*, e001683. [[CrossRef](#)]
103. Sicko, R.J.; Browne, M.L.; Rigler, S.L.; Druschel, C.M.; Liu, G.; Fan, R.; Romitti, P.A.; Caggana, M.; Kay, D.M.; Brody, L.C.; et al. Genetic Variants in Isolated Ebstein Anomaly Implicated in Myocardial Development Pathways. *PLoS ONE* **2016**, *11*, e0165174. [[CrossRef](#)]
104. Hoedemaekers, Y.M.; Caliskan, K.; Majoor-Krakauer, D.; van de Laar, I.; Michels, M.; Witsenburg, M.; ten Cate, F.J.; Simoons, M.L.; Dooijes, D. Cardiac beta-myosin heavy chain defects in two families with non-compaction cardiomyopathy: Linking non-compaction to hypertrophic, restrictive, and dilated cardiomyopathies. *Eur Heart J.* **2007**, *28*, 2732–2737. [[CrossRef](#)]
105. Stampfli, S.F.; Donati, T.G.; Hellermann, J.; Anwer, S.; Erhart, L.; Gruner, C.; Kaufmann, B.A.; Gencer, B.; Haager, P.K.; Muller, H.; et al. Right ventricle and outcome in left ventricular non-compaction cardiomyopathy. *J. Cardiol.* **2020**, *75*, 20–26. [[CrossRef](#)]
106. Moorman, A.; Webb, S.; Brown, N.A.; Lamers, W.; Anderson, R.H. Development of the heart: (1) Formation of the cardiac chambers and arterial trunks. *Heart* **2003**, *89*, 806–814. [[CrossRef](#)]
107. Sylva, M.; van den Hoff, M.J.; Moorman, A.F. Development of the human heart. *Am. J. Med. Genet. A* **2014**, *164*, 1347–1371. [[CrossRef](#)]
108. Hinton, R.B.; Yutzey, K.E. Heart valve structure and function in development and disease. *Annu. Rev. Physiol.* **2011**, *73*, 29–46. [[CrossRef](#)]
109. Combs, M.D.; Yutzey, K.E. Heart valve development: Regulatory networks in development and disease. *Circ. Res.* **2009**, *105*, 408–421. [[CrossRef](#)]
110. Krishnan, A.; Samtani, R.; Dhanantwari, P.; Lee, E.; Yamada, S.; Shiota, K.; Donofrio, M.T.; Leatherbury, L.; Lo, C.W. A detailed comparison of mouse and human cardiac development. *Pediatr. Res.* **2014**, *76*, 500–507. [[CrossRef](#)]

111. Anderson, R.H.; Webb, S.; Brown, N.A.; Lamers, W.; Moorman, A. Development of the heart: (2) Septation of the atriums and ventricles. *Heart* **2003**, *89*, 949–958. [[CrossRef](#)]
112. Perez-Pomares, J.M.; de la Pompa, J.L. Signaling during epicardium and coronary vessel development. *Circ. Res.* **2011**, *109*, 1429–1442. [[CrossRef](#)]
113. Gong, H.; Lyu, X.; Wang, Q.; Hu, M.; Zhang, X. Endothelial to mesenchymal transition in the cardiovascular system. *Life Sci.* **2017**, *184*, 95–102. [[CrossRef](#)]
114. Lincoln, J.; Yutzey, K.E. Molecular and developmental mechanisms of congenital heart valve disease. *Birth Defects Res. A Clin. Mol. Teratol.* **2011**, *91*, 526–534. [[CrossRef](#)]
115. de Vlaming, A.; Sauls, K.; Hajdu, Z.; Visconti, R.P.; Mehesz, A.N.; Levine, R.A.; Slaugenhaupt, S.A.; Hagege, A.; Chester, A.H.; Markwald, R.R.; et al. Atrioventricular valve development: New perspectives on an old theme. *Differentiation* **2012**, *84*, 103–116. [[CrossRef](#)]
116. de Lange, F.J.; Moorman, A.F.; Anderson, R.H.; Manner, J.; Soufan, A.T.; de Gier-de Vries, C.; Schneider, M.D.; Webb, S.; van den Hoff, M.J.; Christoffels, V.M. Lineage and morphogenetic analysis of the cardiac valves. *Circ. Res.* **2004**, *95*, 645–654. [[CrossRef](#)]
117. Lincoln, J.; Alfieri, C.M.; Yutzey, K.E. Development of heart valve leaflets and supporting apparatus in chicken and mouse embryos. *Dev. Dyn.* **2004**, *230*, 239–250. [[CrossRef](#)]
118. Oechslin, E.; Jenni, R. Left ventricular non-compaction revisited: A distinct phenotype with genetic heterogeneity? *Eur. Heart J.* **2011**, *32*, 1446–1456. [[CrossRef](#)]
119. Sedmera, D.; Pexieder, T.; Vuillemin, M.; Thompson, R.P.; Anderson, R.H. Developmental patterning of the myocardium. *Anat. Rec.* **2000**, *258*, 319–337. [[CrossRef](#)]
120. Jimenez-Amilburu, V.; Rasouli, S.J.; Staudt, D.W.; Nakajima, H.; Chiba, A.; Mochizuki, N.; Stainier, D.Y.R. In Vivo Visualization of Cardiomyocyte Apicobasal Polarity Reveals Epithelial to Mesenchymal-like Transition during Cardiac Trabeculation. *Cell Rep.* **2016**, *17*, 2687–2699. [[CrossRef](#)]