Supplementary Information

Selective Inhibition of Human Monoamine oxidase B by 5-Hydroxy-2-Methyl-Chroman-4-One Isolated from an Endogenous Lichen Fungus *Daldinia fissa*

Geum Seok Jeong ¹, Myung-Gyun Kang ², Sang-Ah Han ³, Ji-In Noh ¹, Jong Eun Park ¹, Sang-Jip Nam ⁴, Daeui Park ², Sung-Tae Yee ¹ and Hoon Kim ^{1,*}

Table of Contents

Figure S1. Inhibitory activity of ELF 195 extracts against MAO-A at 20 μ g/mLS3
Figure S2 . Inhibitory activity of ELF 195 extracts against MAO-B at 20 μg/mLS4
Figure S3 . Inhibitory activity of ELF 195 extracts against AChE at 50 μg/mLS5
Figure S4 . Inhibitory activity of ELF 195 extracts against BChE at 50 μg/mLS6
Figure S5 . Antioxidant activity of ELF 195 extract using DPPH at 100 μg/mLS7
Figure S6 . ¹ H NMR spectrum of compound C2 in MeOD- <i>d</i> ₄ S8
Figure S7. ¹³ C NMR spectrum of compound C2 in MeOD- <i>d</i> ₄
Figure S8. HMBC NMR spectrum of compound C2 in MeOD- <i>d</i> ₄ S10
Figure S9. LC/MS chromatogram of compound C2S11
Figure S10. Plots of root mean square deviation during 1 ns MD simulation of MAO-
B in complexes with (<i>R</i>)- and (<i>S</i>)-HMCS12
Figure S11. Plots of root mean square deviation during 1 ns MD simulation of MAO-
A in complexes with (<i>R</i>)- and (<i>S</i>)-HMC



Figure S1. Inhibitory activity of ELF 195 extracts against MAO-A at 20 μ g/mL.



Figure S2. Inhibitory activity of ELF 195 extracts against MAO-B at 20 μ g/mL.



Figure S3. Inhibitory activity of ELF 195 extracts against AChE at 50 μ g/mL.



Figure S4. Inhibitory activity of ELF 195 extracts against BChE at 50 μ g/mL.



Figure S5. Antioxidant activity of ELF 195 extract using DPPH at 100 μ g/mL.



Figure S6. ¹H NMR spectrum of compound **C2**.



Figure S7. ¹³C NMR spectrum of compound **C2**.



Figure S8. HMBC NMR spectrum of compound **C2**.



Figure S9. LC/MS chromatogram of compound C2.



Figure S10. Plots of root mean square deviation during 1000 ps MD simulation of MAO-B in complexes with (*R*)- and (*S*)-HMC.



Figure S11. Plots of root mean square deviation during 1000 ps MD simulation of MAO-A in complexes with (*R*)- and (*S*)-HMC.