

Supplementary materials

Calcium affects polyphosphate and lipid accumulation in Mucoromycota fungi

Simona Dzurendova^{1*}, Boris Zimmermann¹, Achim Kohler¹, Kasper Reitzel², Ulla Gro Nielsen³, Benjamin Xavier Dupuy-Galet¹, Shaun Leivers⁴, Svein Jarle Horn⁴, Volha Shapaval¹

¹Norwegian University of Life Sciences, Faculty of Science and Technology, Drøbakveien 31, 1433 Ås, Norway

²Department of Biology, University of Southern Denmark, Campusvej 55, DK-5230, Odense M, Denmark

³ Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Campusvej 55, DK-5230, Odense M, Denmark

⁴Norwegian University of Life Sciences, Faculty of Chemistry, Biotechnology and Food Science, Christian Magnus Falsens vei 1, 1433 Ås, Norway

* Correspondence: simona.dzurendova@gmail.com; simona.dzurendova@nmbu.no




Figure S1: pH of culture supernatants at the end of the cultivation.




Figure S2: The total concentrations (in g/L) of fungal biomass, lipids and phosphates.




Figure S3: The fatty acid profiles of FAMEs extracted from fungal biomass. Fatty acids which were present in amount higher than 1% are shown, the rest is summed up into ‘others’.