



Supplementary Material

**Table S1.** Univariate analysis between dependant variables and the characteristic of the patients.

	<b>Stress p.trend <sup>1</sup></b>	<b>Anxiety depression p.trend <sup>1</sup></b>	<b>Pain Discomfort p.trend <sup>1</sup></b>	<b>Selfcare p.trend <sup>1</sup></b>	<b>Usual activities p.trend <sup>1</sup></b>	<b>Mobility p.trend <sup>1</sup></b>
age	0.256	0.120	0.017	0.048	<0.001	0.001
Gender	0.575	0.926	0.865	0.556	0.834	0.846
Matrimonial status	0.210	0.627	0.136	0.137	0.686	0.325
residence	0.116	0.099	0.174	0.381	0.687	0.422
Education	0.391	0.027	0.073	0.963	0.008	0.153
Occupation	0.341	0.885	0.333	0.703	0.206	0.638
Household num- ber/size	<0.001	<0.001	0.005	0.063	0.069	0.005
Tuberculosis type	0.924	0.155	0.179	0.183	0.700	0.408
Disease localisation	0.008	0.344	0.517	0.322	0.034	0.045
Treatment history	0.656	0.347	0.449	0.622	0.041	0.148
TB diagnostic	0.005	0.113	0.067	0.227	0.072	0.710
Treatment type	0.852	0.198	0.225	0.324	0.861	0.268
Type multi-drug treatment	0.881	0.267	0.072	0.379	0.761	0.326
Anti-retroviral treatment	0.056	0.001	0.007	0.768	0.063	0.071
Distance site Patient residence	0.001	<0.001	0.182	0.025	0.185	0.313
Length time for drug procurement (in days)	0.004	<0.001	0.129	0.242	0.009	0.004
Treatment initiation	0.383	0.430	0.700	0.991	0.975	0.460
Duration of treatment	0.381	0.202	0.276	0.199	0.019	0.103

<sup>1</sup> p trend computed with the levels (no problem, moderate, serious) for each dependent variable.