

Supplementary Materials: Letermovir for Cytomegalovirus Prophylaxis in Lung Transplant Patients with Valganciclovir-Induced Leukopenia

Arindam Singha, Pamela K. Burcham, April Logan, Zeinab El Boghdadly, Molly M. Howsare, David R. Nunley, Mark E. Lustberg and Brian C. Keller *

Table S1. Detailed patient demographics.

| Patient | Age | Sex | Tx indication | Tx type | CMV status | Leukopenia episodes prior to LET | Median leukopenia duration (d) | WBC nadir (K/uL) | Time to leukopenia resolution after LET (d) | Indication for CMV prophylaxis w/ LET | LET start (POD) |
|---------|-----|-----|---------------|---------|------------|----------------------------------|--------------------------------|------------------|---------------------------------------------|---------------------------------------|-----------------|
| #1 | 27 | M | CF | BLT | D+/R- | 4 | 7 | 0.61 | 44 | Primary | 102 |
| #2 | 65 | F | CF | BLT | D+/R- | 5 | 11 | 0.62 | 9 | Secondary | 182 |
| #3 | 62 | F | IPF | BLT | D+/R- | 6 | 21 | 1.17 | 8 | Secondary | 279 |
| #4 | 54 | F | IPF | BLT | D+/R+ | 4 | 8 | 1.03 | 6 | Primary | 99 |
| #5 | 63 | M | IPF | BLT | D+/R- | 2 | 92.5 | 0.94 | 33 | Primary | 294 |
| #6 | 65 | F | IPF | SLT | D+/R- | 7 | 23 | 1.24 | 38 | Secondary | 767 |
| #7 | 49 | M | CF | BLT | D+/R- | 2 | 58 | 1.10 | 10 | Primary | 321 |
| #8 | 62 | F | COPD | BLT | D+/R- | 4 | 29.5 | 1.48 | 7 | Secondary | 601 |
| #9 | 56 | F | COPD | BLT | D-/R+ | 4 | 23 | 1.11 | 14 | Secondary | 219 |
| #10 | 68 | M | IPF | BLT | D+/R- | 4 | 13 | 2.00 | 8 | Primary | 185 |
| #11 | 68 | M | IPF | BLT | D-/R+ | 1 | 29 | 1.10 | 13 | Primary | 77 |
| #12 | 64 | M | COPD | BLT | D+/R- | 4 | 5 | <0.30 | 5 | Secondary | 118 |
| #13 | 60 | F | COPD | SLT | D+/R- | 2 | 136 | 0.90 | 56 | Secondary | 515 |
| #14 | 60 | M | CLAD | BLT | D-/R+ | 1 | 34 | 2.19 | 7 | Primary | 101 |
| #15 | 27 | M | CF | BLT | D-/R+ | 1 | 112 | 0.90 | 34 | Primary | 154 |
| #16 | 62 | M | IPF | BLT | D+/R+ | 2 | 16.5 | 0.59 | -6* | Primary | 110 |
| #17 | 40 | M | Sarcoidosis | BLT | D+/R- | 1 | 184 | 1.40 | 77 | Secondary | 263 |

Leukopenia episodes were defined as WBC <4 K/uL on consecutive blood samples collected at least 1 day apart. Resolution of leukopenia was defined as WBC ≥ 4K/uL on two consecutive blood samples collected at least 1 day apart. BLT, bilateral lung transplant; CF, cystic fibrosis; CLAD, chronic lung allograft dysfunction; COPD, chronic obstructive pulmonary disease; IPF, idiopathic pulmonary fibrosis; LET, letermovir; LT, lung transplant; POD, post-operative day; SLT, single lung transplant; Tx, transplant. * = Patient #16 had resolution of leukopenia 6 days prior to starting letermovir.



Table S2. Repeated Measures Mixed Model.



| Description | Parameter | Parameter Estimate | p-value |
|-------------|--------------------------------------------|--------------------|---------|
| Intercepts | Valganciclovir | 9.20 | 0.0449 |
| | Letermovir | 7.51 | |
| Slopes | Days Post-Transplant for No Letermovir Yet | -0.0113 | <0.0001 |
| | Days Post-Transplant for Letermovir | -0.0043 | |



Table S3. Laboratory values and filgrastim doses measured or administered before and after letermovir.



| Patient # | WBC Pre (K/uL) | WBC Post (K/uL) | SCr Pre (mg/dL) | SCr Post (mg/dL) | Tacrolimus Pre (ng/mL) | Tacrolimus Post (ng/mL) | Filgrastim doses Pre | Filgrastim doses Post |
|-----------|-------------------|--------------------|--------------------|---------------------|---------------------------|----------------------------|-------------------------|--------------------------|
| 1 | 3.43 | 3.05 | 1.55 | 2.30 | 8.20 | 9.20 | 8 | 2 |
| 2 | 2.86 | 4.24 | 0.98 | 0.98 | 8.00 | 7.90 | 3 | 0 |
| 3 | 3.06 | 4.85 | 2.15 | 1.85 | 5.40 | 4.60 | 1 | 0 |
| 4 | 2.61 | 4.76 | 0.62 | 0.65 | 10.45 | 10.40 | 11 | 0 |
| 5 | 2.00 | 4.60 | 4.40 | 3.57 | 9.70 | 5.25 | 6 | 0 |
| 6 | 4.70 | 6.05 | 1.50 | 2.45 | 7.80 | 14.45 | 3 | 0 |
| 7 | 2.20 | 7.55 | 1.25 | 1.20 | 10.18 | 7.70 | 2 | 0 |
| 8 | 5.00 | 7.71 | 1.51 | 1.35 | 6.70 | 9.60 | 5 | 0 |
| 9 | 1.60 | 2.90 | 2.29 | 1.90 | 11.90 | 5.00 | 17 | 0 |
| 10 | 4.24 | 8.47 | 1.66 | 1.65 | 8.30 | 9.10 | 0 | 0 |
| 11 | 4.60 | 4.48 | 1.00 | 0.96 | 11.35 | 11.40 | 0 | 2 |
| 12 | 4.80 | 7.77 | 1.12 | 0.97 | 7.60 | 8.60 | 9 | 7 |
| 13 | 2.80 | 4.50 | 1.48 | 1.21 | 7.50 | 3.50 | 6 | 0 |
| 14 | 4.59 | 5.91 | 2.28 | 3.60 | 7.65 | 7.80 | 0 | 2 |
| 15 | 1.19 | 5.20 | 1.10 | 1.25 | 12.00 | 9.55 | 4 | 0 |
| 16 | 2.62 | 5.57 | 1.16 | 1.27 | 8.80 | 10.60 | 8 | 0 |
| 17 | 2.42 | 3.09 | 1.00 | 1.22 | 8.50 | 15.20 | 0 | 0 |



WBC significantly increased following letermovir initiation (3.15 K/uL pre-letermovir vs. 5.46 K/uL post-letermovir, $p < 0.0001$). There was no significant change in SCr or tacrolimus trough level after starting letermovir ($p = 0.56$ and 0.48 , respectively). Mean number of filgrastim administrations required to maintain normal WBC levels were significantly lower following initiation of letermovir (4.88 doses pre-letermovir vs. 0.76 doses post-letermovir, $p = 0.003$). SCr, serum creatinine.


























Figure S1. Patient-specific plots of CMV DNAemia from time of transplant through end of study period.