

Selection and Validation of Reliable Reference Genes for qRT-PCR Normalization of *Bursaphelenchus xylophilus* from Different Temperature Conditions and Developmental Stages

Yajing Wu ^{1,2,3,†}, Jiao Zhou ^{2,†}, Yaning Liu ², Yutong Gu ², Hongxia Zhang ², Faheem Ahmad ⁴, Guochang Wang ^{3,*} and Lili Ren ^{1,5*}

¹ Science and Technology Research Center of China Customs, Beijing 100026, China; wyjds@163.com

² State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China; zhoujiao@ioz.ac.cn (J.Z.); liuyanling6206@126.com (Y.L.); gyt715166463@163.com (Y.G.); zhanghx@ioz.ac.cn (H.Z.)

³ School of Resources and Environment, Henan Institute of Science and Technology, Xinxiang 453003, China

⁴ Department of Biosciences, COMSATS University Islamabad (CUI), Park Road, Tarlai Kalan 45550 ICT, Islamabad, Pakistan; faheem.ahmad@comsats.edu.pk

⁵ Institute of Inspection Technology and Equipment, Chinese Academy of Inspection and Quarantine, Beijing 100029, China;

* Correspondence: wgchslbh@163.com (G.W.); renll2021@163.com (L.R.)

† These authors have contributed equally to this work.




Figure S1. Standard curves of eight candidate reference genes. Line relations and determination coefficient (R^2) between Ct values and Log quantity were calculated by SPSS 22.0 software.