

Table S1. Food finding test in 3xTg-AD and NTg mice under naturalistic and forced isolation.

Naturalistic Social Isolation [14] 12-month-old with overnight food deprivation

	NTg Male (n=24)		3xTg-AD Male (n=15)		3xTg AD-ISO Male (n=7)	
	Mean	± SEM	Mean	± SEM	Mean	± SEM
Food finding test						
Lat of sniffing (s)	251	± 63.8	135.7	± 16.2	85.1	± 14.5
Lat of finding food (s)	424.2	± 57.9	152.4	± 15.7	111.8	± 17.1
Lat of eating (s)	912.4	± 121	193	± 17.7	138.1	± 23.6

Forced Social Isolation (current work) – 16-month-old without overnight food deprivation

	NTg Male (n=6)		NTg-ISO Male (n=8)		3xTg-AD Male (n=10)		3xTg AD-ISO Male (n=10)	
	Mean	± SEM	Mean	± SEM	Mean	± SEM	Mean	± SEM
Food finding test								
Lat of sniffing (s)	436.5	± 50.8	967.3	± 212	919.4	± 102.5	491.7	± 100.1
Lat of finding food (s)	516.9	± 60.1	1349.5	± 305.6	973.7	± 103.3	578.5	± 124.3
Lat of eating (s)	1423.5	± 258.4	2106.5	± 295.2	2693.7	± 63.9	47.5	± 3.5