

| Characteristic | knowledge | | | Attitudes | | | | |
|-----------------------|---------------------|---------------------------------------|------------------------------|--|--|---|-------------------------------------|-------------------|
| | Typology OR (CI) | Countries of prevalence OR (CI) | Cases detected OR (CI) | Discuss in the outpatient clinic OR (CI) | Asked about other daughters OR (CI) | Asked other professionals OR (CI) | Report to authorities OR (CI) | Ignore OR (CI) |
| Age | | | | | | | | |
| 20-40 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 41-50 | 1.5 (0.9-2.5) | 0.8 (0.6-1.2) | 0.6 (0.3-1.3) | 0.7 (0.2-2.4) | 2.1 (0.5-9.2) | 0.9 (0.3-2.9) | 0 | 0.5 (0.2-1.8) |
| >50 | 1.4 (0.8-2.3) | 0.8 (0.5-1.1) | 0.8 (0.4-1.5) | 1.0 (0.3-3.3) | 2.5 (0.5-11.6) | 0.3 (0.1-1.7) | 0 | 0.8 (0.3-2.4) |
| Profession | | | | | | | | |
| Nursing | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Gynaecologist | 8.5 (2.5-29.1)* | 0.36 (0.04-2.8) | 14.7 (3.5-61.6) | 21.9 (2.0-244.1)* | 14.9 (1.4-160.7)* | 0 | 0 | 39.5 (6.2-250.9) |
| Family medicine | 1.0 (0.7-1.7) | 0.84 (0.6-1.2) | 2.0 (1.0-4.3)* | 3.7 (0.8-16.0) | 0.5 (0.5-4.3) | 0.6 (0.1-5.8) | 0 | 3.0 (0.9-113.7) |
| Midwife | 4.3(2.1-8.8)* | 1.7 (0.9-3.4) | 48.5 (21.3-110.1)* | 43.7 (10.7-178.4)* | 28.9 (7.6-110.7)* | 83.2 (19.6-352.5)* | 0 | 32.0 (9.0-113.2) |
| Obstetrician | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Paediatrician | 1.5(0.7-3.0) | 1.9 (1.4-3.2)* | 3.0 (1.0-8.5) | 2.7 (0.3-26.8) | 1.9 (0.2-17.4) | 3.2 (0.3-31.9) | 0 | 0 |
| Gender | | | | | | | | |
| Male | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Female | 0.7(0.4-1.1) | 0.9 (0.6-1.3) | 0.8 (0.4-1.7) | 0.6 (0.2-2.2) | 0.8 (0.2-3.3) | 3.0 (0.8-11.4) | 0 | 0.4 (0.1-1.5) |
| Training | | | | | | | | |
| Training not received | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Training received | 3.0 (1.9-4.7)* | 2.3 (1.6-3.4)* | 1.0 (0.5-2.1) | 2.5 (0.9-7.0) | 1.9 (0.6-6.3) | 1.2 (0.4-4.2) | 0 | 0.5 (0.1-1.9) |

Results are expressed using Odds Ratio and CI 95%

* p < 0.05 on comparison with the reference category