



Supplementary Table S1. Unadjusted and adjusted odds ratios for the association between basic risk factors on gestational hypertension (GH) and preeclampsia (PE) (set of data).

Risk factors	Cases/Controls	OR (95% CI), <i>p</i> GH risk	AOR-a (95% CI), <i>p</i>
Maternal age (years):			
≥40	16/59	3.23 (1.41–7.38); 0.005	2.97 (1.2–7.32) 0.018
Age 18–24	2/36	0.66 (0.14–3.12); 0.602	0.51 (0.1–2.53) 0.413
25–29 years	11/131	1	1
Pre-pregnancy BMI (kg/m^2):			
Obesity (≥ 30)	31/58	5.60 (3.32–9.43); <0.001	4.72 (2.73–8.14) <0.001
Overweight (25–29.9)	30/139	2.26 (1.39–3.68); 0.001	2.04 (1.23–3.37) 0.005
Underweight (< 18.5)	1/44	0.24 (0.03–1.76); 0.160	0.23 (0.03–1.69) 0.148
Normal BMI (18.5–24.9)	51/534	1	1
GWG above the range			
GWG in the range	62/263	2.45 (1.53–3.92); < 0.001	1.9 (1.15–3.12); 0.012
GWG below the range	29/301	1	1
Smoking in the 1st tr.	22/211	1.08 (0.61–1.94); 0.790	1.09 (0.59–2.01); 0.779
Smoking before pregnancy	17/37	3.60 (1.94–6.68); <0.001	3.51 (1.79–6.89); <0.001
Never smoked	31/133	1.83 (1.16–2.87); 0.009	1.64 (1.01–2.66); 0.044
Prior GH/PE	82/642	1	1
No prior GH/PE	12/4	22.9 (7.25–72.36); <0.001	37.98 (11.16–129.26); <0.001
Primiparity	101/771	1	1
Multiparity	53/318	1.27 (0.85–1.89); 0.238	1.90 (1.22–2.96); 0.005
Infertility treatment	60/457	1	1
No Infertility treatment	8/29	1.96 (0.87–4.4); 0.103	1.65 (0.68–4.01); 0.269
PE risk			
Maternal age (years):			
≥40	1/59	1.11 (0.1–12.49); 0.933	0.94 (0.08–11.61); 0.963
25–29	2/131	1	1
18–24	2/36	3.64 (0.5–26.74); 0.204	3.26 (0.42–25.04); 0.256
Pre-pregnancy BMI (kg/m^2):			
Obesity (≥ 30)	9/58	9.21 (3.52–24.11); < 0.001	8.68 (3.25–23.18); <0.001
Overweight (25–29.9)	4/139	1.71 (0.52–5.63); 0.379	1.68 (0.5–5.63); 0.397
Underweight (< 18.5)	2/44	2.70 (0.57–12.87); 0.213	2.56 (0.52–12.62); 0.248
Normal BMI (18.5–24.9)	9/534	1	1
GWG above the range			
GWG in the range	11/263	1.57 (0.62–3.97); 0.337	1.15 (0.44–3); 0.770
GWG below the range	8/301	1	1
Smoking in the 1st tr.	5/211	0.89 (0.29–2.76); 0.842	0.84 (0.26–2.67); 0.768
Smoking before pregnancy	3/37	2.60 (0.74–9.16); 0.136	2.79 (0.76–10.27); 0.124
Never smoked	4/133	0.97 (0.33–2.87); 0.950	0.88 (0.29–2.65); 0.817
Prior GH/PE	20/642	1	1
No prior GH/PE	3/4	27.54 (5.8–130.83); <0.001	31.11 (5.83–165.98); <0.001
Primiparity	21/771	1	1
Multiparity	11/318	1.22 (0.54–2.75); 0.638	1.55 (0.64–3.77); 0.332
Infertility treatment	13/457	1	1
No infertility treatment	3/29	3.68 (1.04–13.03); 0.044	4.29 (1.06–17.39); 0.041
	21/746	1	1

* AOR-a: adjusted odds ratios (and 95% confidence intervals) of GH and PE calculated in multiple logistic regression in model-a; the results were adjusted for maternal age, pre-pregnancy BMI, primiparity, gestational weight gain (GWG) out of the range and smoking in the first trimester (*p*-value < 0.05 was assumed to be significant). GH: gestational hypertension; PE: preeclampsia. Controls: normotensive women.

Supplementary Table S2. Unadjusted and adjusted odds ratios for the association between paternal and maternal hypertension on gestational hypertension (GH) and preeclampsia (PE), compared to ‘Absence of hypertension in the parents’

Risk factors/Hypertension in the parents	Cases/Controls	OR (95% CI); <i>p</i>	AOR-a (95% CI); <i>p</i> *	AOR-b (95% CI); <i>p</i> *
GH risk				
In the mother	31/137	2.01 (1.25–3.21); 0.004	1.61 (0.97–2.67); 0.065	1.46 (0.86–2.48); 0.165
In the father	33/135	2.17 (1.36–3.45); 0.001	2.09 (1.27–3.44); 0.004	2.03 (1.21–3.42); 0.008
In the mother or father	51/230	1.97 (1.31–2.94); 0.001	1.75 (1.14–2.69); 0.010	1.65 (1.06–2.58); 0.028
In the mother and father **	13/42	2.75 (1.40–5.40); 0.003	2.31 (1.11–4.82); 0.025	2.14 (0.98–4.65); 0.056
Ref ***	61/541	1	1	1
PE risk				
In the mother	11/137	3.95 (1.68–9.30); 0.002	3.28 (1.34–8.04); 0.009	3.12 (1.24–7.85); 0.016
In the father	5/135	1.82 (0.62–5.33); 0.274	1.58 (0.52–4.76); 0.416	1.37 (0.42–4.53); 0.605
In the mother or father	13/230	2.78 (1.23–6.30); 0.014	2.35 (1.02–5.45); 0.046	2.15 (0.89–5.17); 0.088
In the mother and father **	3/42	3.51 (0.94–13.08); 0.061	2.78 (0.69–11.24); 0.151	2.57 (0.59–11.2); 0.207
Ref ***	11/541	1	1	1

* AOR: adjusted odds ratios (and 95% confidence intervals) calculated in multiple logistic regression after adjusted for maternal age, pre-pregnancy BMI, primiparity, gestational weight gain (GWG) out of the range and smoking in the first trimester (AOR-a) plus prior hypertension in pregnancy and infertility treatment (AOR-b) (*p*-value < 0.05 was assumed to be significant); ** In the mother and father simultaneously; *** Reference category: ‘Absence of hypertension in the parents’. Cases: GH i.e. gestational hypertension; PE i.e. preeclampsia; Controls: normotensive women.