

Table S2. Primary outcome in the Quit with US trial by separating daily and nondaily smokers

Exhaled CO concentration level verified 7-day point prevalence abstinence ¹	Daily smokers				Nondaily smokers			
	Quit with US, <i>n</i> (%)	Control, <i>n</i> (%)	RR (95% CI)	<i>p</i>	Quit with US, <i>n</i> (%)	Control, <i>n</i> (%)	RR (95% CI)	<i>p</i>
Intention-to-treat								
analysis	(<i>n</i> = 75)	(<i>n</i> = 83)			(<i>n</i> = 62)	(<i>n</i> = 53)		
Abstainers	28 (37.3)	8 (9.6)	3.87 (1.88 to 7.96)	<0.001	52 (83.9)	34 (64.2)	1.31 (1.04 to 1.64)	0.018
Nonabstainers	47 (62.7)	75 (90.4)	1.00		10 (16.1)	19 (35.8)	1.00	
Per protocol analysis	(<i>n</i> = 64)	(<i>n</i> = 74)			(<i>n</i> = 55)	(<i>n</i> = 45)		
Abstainers	28 (43.8)	8 (10.8)	4.05 (1.99 to 8.24)	<0.001	52 (94.6)	34 (75.6)	1.25 (1.05 to 1.49)	0.009
Nonabstainers	36 (56.2)	66 (89.2)	1.00		3 (5.4)	11 (24.4)	1.00	

RR, risk ratio; CI, confidence intervals; ¹ Measured by a self-report of continuous abstinence from smoking in the previous 7 consecutive days plus an exhaled CO concentration level of ≤6 ppm.