

Supplementary Materials

Relationship between antioxidant activity and ligand basicity in the dipicolinate series of oxovanadium(IV) and dioxovanadium(V) complexes

Joanna Drzeżdżon¹, Marta Pawlak¹, Natalia Matyka¹, Artur Sikorski¹,
Barbara Gawdzik^{2,*} and Dagmara Jacewicz¹

¹ Faculty of Chemistry, University of Gdańsk, Wita Stwosza 63, 80-308 Gdańsk, Poland;
joanna.drzezdzon@ug.edu.pl (J.D.); marta.pawlak0812@gmail.com (M.P.); natmatyla@gmail.com (N.M.);
artur.sikorski@ug.edu.pl (A.S.); dagmara.jacewicz@ug.edu.pl (D.J.);

² Institute of Chemistry, Jan Kochanowski University, Świętokrzyska 15 G, 25-406 Kielce, Poland;
b.gawdzik@ujk.edu.pl

* Correspondence: b.gawdzik@ujk.edu.pl; Tel.: +48 41 349 70 16




Figure S1. IR spectrum of $[\text{VO}(\text{dipic})(\text{H}_2\text{O})_2] \cdot 2 \text{H}_2\text{O}$.




Figure S2. IR spectrum of $[\text{VO}(\text{dipic})(\text{phen})] \cdot 3 \text{H}_2\text{O}$.




Figure S3. IR spectrum of [VO(dipic)(bipy)] · H₂O.




Figure S4. MALDI-TOF-MS spectrum of $[\text{VO}(\text{dipic})(\text{H}_2\text{O})_2] \cdot 2 \text{H}_2\text{O}$ (CCA matrix).




Figure S5. MALDI-TOF-MS spectrum of [VO(dipic)(phen)] · 3 H₂O (CCA matrix).




Figure S6. MALDI-TOF-MS spectrum of [VO(dipic)(bipy)] · H₂O (CCA matrix).




Figure S7. MALDI-TOF-MS spectrum of $[\text{VO}(\text{dipic})(\text{H}_2\text{O})_2] \cdot 2 \text{H}_2\text{O}$ (DHB matrix).




Figure S8. MALDI-TOF-MS spectrum of [VO(dipic)(phen)] · 3 H₂O (DHB matrix).




Figure S9. MALDI-TOF-MS spectrum of [VO(dipic)(bipy)] · H₂O (DHB matrix).




Figure S10. ^1H NMR spectrum of $[\text{VO}(\text{dipic})(\text{bipy})] \cdot \text{H}_2\text{O}$.




Figure S11. ^{13}C NMR spectrum of $[\text{VO}(\text{dipic})(\text{bipy})] \cdot \text{H}_2\text{O}$.