

Supplementary Material

Marrubiin Inhibits Peritoneal Inflammatory Response Induced by Carrageenan Application in C57 Mice

Niko S. Radulović ^{1,*}, Miljana R. Đorđević Zlatković ¹, Nikola M. Stojanović ², Milan S. Nešić ¹, Dragan B. Zlatković ¹, Milena S. Potić Floranović ³, Dragana S. Tričković Vukić ³ and Pavle J. Randjelović ²

¹ Department of Chemistry, Faculty of Sciences and Mathematics, University of Niš, 18000 Niš, Serbia; miljana.djordjevic@pmf.edu.rs (M.R.Đ.Z.); nesicmilanvl@gmail.com (M.S.N.); dragan.zlatkovic@gmail.com (D.B.Z.)

² Department of Physiology, Faculty of Medicine, University of Niš, 18000 Niš, Serbia; nikola.st90@yahoo.com (N.M.S.); pavleus@gmail.com (P.J.R.)

³ Scientific Research Centre for Biomedicine, Faculty of Medicine, University of Niš, 18000 Niš, Serbia; milenapotic@yahoo.com (M.S.P.F.); draganavukic18@gmail.com (D.S.T.V.)

* Correspondence: nikoradulovic@yahoo.com

Table of contents:

| | page |
|---|------|
| Figure S1: ¹ H NMR spectrum of marrubiin (MAR) | 2 |
| Figure S2: ¹³ C NMR spectrum of MAR | 2 |
| Figure S3: ¹ H NMR spectrum of MAR after 10 years of storage at 4 °C | 3 |




Figure S1. ^1H NMR (400 MHz) spectrum of marrubiin (MAR) in CDCl_3




Figure S2. ^{13}C NMR (100.6 MHz) spectrum of MAR in CDCl_3




Figure S3. ^1H NMR (400 MHz, CDCl_3) spectrum of MAR after 10 years of storage at 4 °C. No degradation of MAR can be observed.