

Article

Ocean acidification induces changes in virus-host relationships in Mediterranean benthic ecosystems

Michael Tangherlini^{1*}, Cinzia Corinaldesi², Francesca Ape³, Silvestro Greco⁴, Teresa Romeo^{5,6}, Franco Andaloro⁵ and Roberto Danovaro⁷

¹ Department of Research Infrastructures for marine biological resources, Stazione Zoologica Anton Dohrn, Fano Marine Centre, Viale Adriatico 1-N, 61032 Fano, Italy; michael.tangherlini@szn.it

² Department of Materials, Environmental Sciences and Urban Planning, Polytechnic University of Marche, Via Brecce Bianche, 60131 Ancona, Italy; c.corinaldesi@staff.univpm.it

³ Institute of Anthropic Impacts and Sustainability in Marine Environment - National Research Council (IAS-CNR), Lungomare Cristoforo Colombo n. 4521 (ex complesso Roosevelt), Località Addaura, 90149 Palermo, Italy; francesca.ape@ias.cnr.it

⁴ Department of Integrated Marine Ecology, Stazione Zoologica Anton Dohrn, Rome; silvestro.greco@szn.it

⁵ Department of Integrated Marine Ecology, Stazione Zoologica Anton Dohrn, Sicily Marine Centre; teresa.romeo@szn.it; franco.andaloro@szn.it.

⁶ National Institute for Environmental Protection and Research, ISPRA, Italy; teresa.romeo@isprambiente.it

⁷ Department of Life and Environmental Sciences, Polytechnic University of Marche, Via Brecce Bianche, 60131 Ancona, Italy; r.danovaro@staff.univpm.it


* Correspondence: michael.tangherlini@szn.it;

Supplementary materials


Supplementary Figures 1-3



Supplementary Figure S1. Map of sampling areas and sampling sites.



Scheme 2. Values of pH for all the sites investigated.



Scheme 3. CLR-transformed relative abundances of differentially abundant taxa classified at domain (A), phylum (B), class (C), order (D) and family (E) levels, with asterisks displaying significant differences between clusters (* for p-values < 0.05, ** for p-values < 0.01, *** for p-values < 0.001).