

Article

Ageritin—The Ribotoxin-like Protein from Poplar Mushroom (*Cyclocybe aegerita*) Sensitizes Primary Glioblastoma Cells to Conventional Temozolomide Chemotherapy

Rossella Rotondo ^{1,†}, Sara Ragucci ^{2,†}, Salvatore Castaldo ¹, Nicola Landi ², Maria Antonietta Oliva ¹, Paolo V. Pedone ², Antimo Di Maro ^{2,*‡} and Antonietta Arcella ^{1,*‡}

¹ IRCCS Istituto Neurologico Mediterraneo NEUROMED, Via Atinense 18, 86077-Pozzilli, IS, Italy; rossella-ross1988@gmail.com (R.R.); castaldosal90@gmail.com (S.C.); mariaantonieta.oliva@neuromed.it (M.A.O.)

² Department of Environmental, Biological and Pharmaceutical Sciences and Technologies (DiSTABiF), University of Campania “Luigi Vanvitelli”, Via Vivaldi 43, 81100-Caserta, Italy; sara.ragucci@unicampania.it (S.R.); nicola.landi@unicampania.it (N.L.); paolovincenzo.pedone@unicampania.it (P.V.P.)

* Correspondence: antimo.dimaro@unicampania.it (A.D.M.); arcella@neuromed.it (A.A.)

† These authors contributed equally to this work.

‡ These authors share equal senior authorship.




Figure S1: Western blot analysis of NF-κB p50 and p65 subunits in long-term Ageritin-treated patient-derived glioblastoma cell lines.