

High-Resolution Bioassay Profiling with Complemented Sensitivity and Resolution for Pancreatic Lipase Inhibitor Screening

Jingyi Jian ^{1,†}, Jiaming Yuan ^{1,†}, Yu Fan ^{1,†}, Jincai Wang ¹, Tingting Zhang ¹, Jeroen Kool ² and Zhengjin Jiang ^{1,*}

¹ Institute of Traditional Chinese Medicine & Natural Products, College of Pharmacy/Guangdong Province Key Laboratory of Pharmacodynamic Constituents of TCM and New Drugs Research/International Cooperative Laboratory of Traditional Chinese Medicine Modernization and Innovative Drug Development of Ministry of Education (MOE) of China, Jinan University, Guangzhou 510632, China

² Division of BioAnalytical Chemistry, Amsterdam Institute of Molecules, Medicines and Systems, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, The Netherlands

* Correspondence: jzjjackson@hotmail.com

† These authors contributed equally to this work.




Figure S1. Dose-response curve of EGCG (IC_{50} of EGCG: $0.97 \mu\text{M}$; R^2 : 0.993).




Figure S2. Dose-response curve of luteolin (IC_{50} of luteolin: $30.57 \mu\text{M}$; R^2 : 0.999).