

## SUPPLEMENTARY MATERIAL

**Table S1:** Number of papers found in the literature search for the period 2000-2020 according to the keyword strings and the databases used, filtered and selected. Note that no article published before 2000 was retrieved using search strings.

Keyword string	ISI WOK			SCOPUS		
	Search	Filtered	Selected	Search	Filtered	Selected
LiDAR* AND biodiversity*	648	317	62	380	216	48
LiDAR* AND vertebrates*	853	23	18	13	12	12
LiDAR* AND mammals*	592	28	25	25	22	16
LiDAR* AND birds*	229	87	75	122	126	68
LiDAR* AND reptiles*	32	4	3	4	3	2
LiDAR* AND amphibians*	5	4	2	5	3	2
LiDAR* AND invertebrates*	254	9	2	13	11	1
LiDAR* AND insects*	138	43	9	104	86	7
LiDAR* AND fungi*	41	10	1	12	9	3
LiDAR* AND lichens*	9	3	3	7	3	3
LiDAR* AND bryophytes*	15	5	5	16	8	5
Total	2816	533	205	701	499	167

**Table S2:** List of species recorded in each of the 173 articles reviewed. #: more than 20 species recorded; N/A: data not available.

<b>Birds</b>			
<b>Order</b>	<b>Common name</b>	<b>Scientific name</b>	<b>Reference</b>
Charadriiformes	Eastern hooded plover	<i>Thinornis cucullatus cucullatus</i>	Ehmke et al. 2016
Charadriiformes	Marbled murrelet	<i>Brachyramphus marmoratus</i>	Hagar et al. 2014
Galliformes	Capercaillie	<i>Tetrao urogallus</i>	Graf et al. 2009, Milanese et al. 2017, Glad et al. 2020
Galliformes	Capercaillie, hazel grouse	<i>Tetrao urogallus, Tetrates bonasia</i>	Vauhkonen & Imponen, 2016, Kortmann et al. 2018(a)
Galliformes	Hazel grouse	<i>Tetrastes bonasia</i>	Bae et al.2014, Zellweger et al. 2014
Galliformes	Greater sage-grouse	<i>Centrocercus urophasianus</i>	Zabihi et al. 2019
Passeriformes	Eurasian sky lark	<i>Alauda arvensis</i>	Mason et al. 2003
Passeriformes	Eurasian sky lark, great tit, Eurasian blue tit	<i>Alauda arvensis, Parusmajor, Cyanistes caeruleus</i>	Bradbury et al. 2005
Passeriformes	Salt-marsh sparrow	<i>Ammodramus caudacutus</i>	Meiman et al. 2012
Passeriformes	Brown creeper	<i>Certhia americana</i>	Vogeler et al. 2013, Vierling et al 2014
Passeriformes	Wood thrush	<i>Hylocichla mustelina</i>	Jirinec et al. 2016
Passeriformes	Red-backed shrike	<i>Lanius collurio</i>	Svendsen et al 2015
Passeriformes	Great tit	<i>Parus major</i>	Hinsley et al. 2006, Hill & Hinsley, 2015
Passeriformes	Great tit, Eurasian blue tit	<i>Parus major, Cyanistes caeruleus</i>	Hinsley et al. 2002, Hinsley et al. 2008, Hinsley et al. 2009(a), Redhead et al 2013
Passeriformes	Siberian jay	<i>Perisoreus infaustus</i>	Klein et al. 2020
Passeriformes	Wood warbler	<i>Phylloscopus sibilatrix</i>	Huber et al 2016, Huber et al. 2017
Passeriformes	Wood warbler	<i>Phylloscopus trochilus</i>	Bellamy et al 2009
Passeriformes	Marsh tit	<i>Poecile palustris</i>	Broughton et al. 2006, Broughton et al. 2012a Broughton et al. 2012b Broughton et al. 2014
Passeriformes	Black-throated blue warbler	<i>Setophaga caerulescens</i>	Goetz et al. 2010
Passeriformes	Cerulean warbler	<i>Setophaga cerulea</i>	Barnes et al. 2006
Passeriformes	Dark-eyed junco, warbling vireo	<i>Junco hyemalis, Vireo gilvus</i>	Clawges et al. 2008
Passeriformes	Eurasian black cap, Common chaffinch	<i>Sylvia atricapilla, Fringilla coelebs</i>	Renner et al. 2016
Passeriformes	Lapland longspur, white-crowned Sparrow	<i>Calcarius lapponicus, Zonotrichia leucophrys</i>	Boelman et al 2016
Passeriformes	Several species	<i>Regulus regulus, Phylloscopus trochilus, Fringilla coelebs, Erithacus rubecula, Parus major,</i>	Eldegard et al. 2016

		<i>Anthus trivialis</i> , <i>Troglodytes troglodytes</i> , <i>Carduelis spinus</i>	
Passeriformes	Several species	<i>Alauda arvensis</i> , <i>Anthus pratensis</i> , <i>A. trivialis</i> , <i>Saxicola rubicola</i> , <i>Turdus philomelos</i> , <i>Lophophanes cristatus</i> , <i>Certhia brachydactyla</i> , <i>Carduelis cannabina</i> , <i>Emberiza citrinella</i>	Ficetola et al. 2014
Passeriformes	Several species	<i>Setophaga striata</i> , <i>S. caerulescens</i> , <i>S. magnolia</i> , <i>S. coronata</i> , <i>Seiurus aurocapilla</i> , <i>Vireo olivaceus</i> , <i>Junco hyemalis</i> , <i>Empidonax flaviventris</i>	Swatantran et al. 2012
Passeriformes	Several species	<i>Setophaga striata</i> , <i>S. caerulescens</i> , <i>S. magnolia</i> , <i>S. coronata</i> , <i>Seiurus aurocapilla</i> , <i>Vireo olivaceus</i> , <i>Junco hyemalis</i> , <i>Empidonax flaviventris</i>	Seavy et al. 2009
Passeriformes	Several species	<i>Hemignathus virens</i> , <i>Himatione sanguinea</i> , <i>Myadestes obscurus</i> , <i>Zosterops japonicus</i> , <i>Garrulax canorus</i>	Boelman et al. 2007
Passeriformes	Several species	#	Müller et al. 2009, Sasaki et al. 2016
Piciformes	Red-cockaded woodpecker	<i>Leuconotopicus borealis</i>	Smart et al. 2012, Trainor et al. 2013, Garabedian et al. 2014, Garabedian et al 2017, Garabedian et al 2018, Garabedian et al. 2020
Piciformes	Lewis's woodpecker	<i>Melanerpes lewis</i>	Vogeler et al. 2016
Piciformes	Black-backed woodpecker	<i>Picoides arcticus</i>	Casas et al. 2016
Piciformes	Three-toed woodpecker	<i>Picoides tridactylus</i>	Zielewska et al. 2018
Piciformes	Red-naped sapsucker	<i>Sphyrapicus nuchalis</i>	Vierling et al. 2013
Strigiformes	Northern spotted owl	<i>Strix occidentalis caurina</i>	Sovern et al. 2019, Hagar et al. 2020
Strigiformes	Northern spotted owl, barred owl	<i>Strix occidentalis caurina</i> , <i>Strix varia</i>	Jenkins et al 2019, Mangan et al. 2019
Strigiformes	California spotted owl	<i>Strix occidentalis occidentalis</i>	Garcia-Feced et al. 2011, North et al. 2017, Gallagher et al. 2019
Several	Several species	<i>Pavo muticus</i> , <i>Treron phayrei</i> , <i>Psittaculaeupatria</i> , <i>P. cularoseata</i> , <i>Ploceus hypoxanthus</i> , <i>Polihierax insignis</i> , <i>Picus rabieri</i>	Singh et al. 2017
Several	Several species	<i>Catharus ustulatus</i> , <i>Geothlypis tolmiei</i> , <i>Oreothlypis celata</i> , <i>Empidonax oberholseri</i> , <i>Leuconotopicus villosus</i> , <i>Colaptes auratus</i> , <i>Sitta canadensis</i> , <i>Poecile gambeli</i>	Lorenz et al. 2015
Several	Several species	<i>Tetrao urogallus</i> , <i>Tetrastes bonasia</i> , <i>Picoides tridactylus</i> , <i>Glaucidium passerinum</i>	Zellweger et al. 2013

Several	Several species	<i>Parus major, Poecile montanus, P. palustris, Sitta europea, Periparus ater, Cyanistes caeruleus, Garrulus glandarius, Lophophane cristatus, Dendrocopos major, Fringilla coelebs, Pyrrhula pyrrhula</i>	Renner et al. 2012
Several	Several species	<i>Laniu scollurio, Crex crex, Otus scops, Picus viridis</i>	Tattoni et al. 2012
Several	Several species	<i>Anous minutus, A. stolidus, Onychoprion lunatus, Gygis alba, Fregata minor, Sula dactylatra, S. sula, Phaethon rubricauda</i>	Hatfield et al. 2012
Several	Several species	<i>Himatione sanguinea, Chlorodrepanis virens, Myadestes obscurus, Drepanis coccinea, Chasiempis sandwichensis, Zosterops japonicus, Haemorhous mexicanus, Leiothrix lutea, Garrulax canorus, Buteo solitarius, Lophura leucomelanos</i>	Flaspohler et al. 2010
Several	Several species	<i>Empidonax oberholseri, Leuconotopicus villosus, Melanerpes lewis, Dryobates pubescens</i>	Martinuzzi et al. 2009
Several	Several species	<i>Anthus trivialis, Fringilla coelebs, Phylloscopus trochilus, Pluvialis apricaria, Tringa totanus, T. glareola, Vanellus vanellus, Anthus pratensis, Alauda arvensis</i>	Leivits&Leivits, 2009
Several	Several species	#	Goetz et al. 2007 Hinsley et al. 2009(b), Müller et al. 2010, Lesak et al. 2011, Jones et al. 2013, Young et al. 2013, Goetz et al 2014, Vogeler et al. 2014, Weisberg et al. 2014, Vihervaara et al. 2015, Wallis et al. 2016, Swift et al. 2017, Bae et al. 2018, Melin et al. 2018, Mononen et al. 2018, Kellner et al. 2018, Adams & Matthews, 2019, Melin et al. 2019 Khanaposhtani et al. 2019, Taillie et al. 2019
Several	Several species	N/A	Davenport et al. 2000, Coops et al. 2016, Carrasco et al. 2019

## Mammals

Chiroptera	Western barbastelle	<i>Barbastella barbastellus</i>	Carr et al. 2018, Kortmann et al. 2018 (b)
Chiroptera	Several species	N/A	Lacki et al 2017
Chiroptera	Several species	<i>Myotis</i> spp., <i>Plecotus</i> spp., <i>Pipistrellus</i> spp., <i>Hypsugos avii</i> , <i>Eptesicus</i> spp., <i>Nyctalus</i> spp., <i>Vespertilio murinus</i>	Froidevaux et al. 2016
Chiroptera	Several species	<i>Pipistrellus</i> spp., <i>Vespertilio</i> spp., <i>Eptesicus</i> spp., <i>Nyctalus</i> spp., <i>Myotis</i> spp.	Müller et al. 2013
Chiroptera	Several species	<i>Pipistrellus pygmaeus</i> , <i>P. nathusii</i> , <i>P. pipistrellus</i> , <i>Myotis</i> sp., <i>Plecotus</i> sp., <i>Barbastella barbastellus</i> , <i>Nyctalus leisleri</i> , <i>N. noctula</i>	Jung et al. 2012
Artiodactyla	White-tailed deer	<i>Odocoileus virginianus</i>	Kroeger et al. 2020
Artiodactyla	Mule deer	<i>Odocoileus hemionus</i>	Coops et al. 2010
Artiodactyla	Several species	<i>Odocoileus</i> spp., <i>Cervus canadensis</i>	Hull et al. 2019
Artiodactyla	Moose	<i>Alces alces</i>	Melin et al. 2013, Melin et al. 2014, Melin et al. 2016
Artiodactyla	European roe deer	<i>Capreolus capreolus</i>	Ewald et al. 2014, Lone et al. 2014
Artiodactyla	Sable antelope	<i>Hippotragus niger</i>	Asner et al 2015
Artiodactyla	Migratory woodland caribou	<i>Rangifer tarandus caribou</i>	Finnegan et al. 2018
Carnivora	Pacific marten	<i>Martes caurina</i>	Tweedy et al. 2019
Carnivora	Fisher	<i>Pekania pennanti</i>	Zhao et al. 2012, Blomdahl et al. 2019
Carnivora	Jaguar	<i>Panthera onca</i>	de Thoisy et al. 2016
Carnivora	Lion	<i>Panthera leo</i>	Loarie et al. 2013, Davies et al 2016(a)
Carnivora	Several species	<i>Lycalopexculpaesus</i> , <i>Lycalopexgriseus</i> , <i>Lycalopexfulvipes</i> , <i>Leopardusguigna</i>	Moreira-Arce et al. 2016
Lagomorpha	Snowshoe hare	<i>Lepusamericanus</i>	Kawaguchi & Desrochers, 2017
Lagomorpha	New England cottontail	<i>Sylvilagustransitionalis</i>	Amaral et al. 2016
Primates	Gray's bald-faced saki	<i>Pithecia irrorata</i>	Palminteri et al. 2012
Primates	Bornean orangutan	<i>Pongo pygmaeus</i>	Davies et al. 2019, Seaman et al. 2019
Primates	Bornean white-bearded gibbon, maroon leaf monkey	<i>Hylobates albibarbis</i> , <i>Presbytis rubicunda</i>	Singh et al. 2018
Primates	Several species	<i>Alouatta palliata</i> , <i>Ateles geoffroyi</i> , <i>Cebus capucinus</i> ,	McLean et al. 2016
Rodentia	Wood mouse	<i>Apodemus sylvaticus</i>	Jaime-Gonzalez et al. 2017
Rodentia	Red tree vole	<i>Arborimus longicaudus</i>	Linnell et al. 2017
Rodentia	Hazel dormouse	<i>Muscardinus avellanarius</i>	Goodwin et al. 2018
Rodentia	Delmarva fox squirrel	<i>Sciurus nigercinereus</i>	Nelson et al. 2005
Rodentia	Red squirrel	<i>Sciurus vulgaris</i>	Flaherty et al. 2014
Rodentia	Douglas squirrel, American red squirrel	<i>Tamiasciurus douglasii</i> , <i>T. hudsonicus</i>	Fotis et al. 2020
Rodentia	American red squirrel	<i>Tamiasciurus hudsonicus</i>	Hatten, 2014

Several	Several species	<i>Blarina brevicauda</i> , <i>Glaucomys</i> sp., <i>Microtus pennsylvanicus</i> , <i>Myodes gapperi</i> , <i>Napaeozapus insignis</i> , <i>Peromyscus leucopus</i> , <i>Peromyscus maniculatus</i> , <i>Sorex arcticus</i> , <i>S. cinereus</i> , <i>S. hoyi</i> , <i>Tamias minimus</i> , <i>T. striatus</i> , <i>T. ciurus hudsonicus</i> , <i>Zapus hudsonius</i>	Schooler et al. 2019
Several	Several species	<i>Panthera leo</i> , <i>Tragelaphus strepsiceros</i> , <i>T. oryx</i> , <i>Syncerus caffer</i> , <i>Equus quagga</i> , <i>Alcelaphus buselaphus</i>	Davies et al. 2016 (b)
<b>Invertebrates</b>			
Blattodea	Termites	<i>Macrotermes</i> spp.	Levick et al 2010, Davies et al. 2014, Davies et al. 2016
Coleoptera	Beetles	<i>Agonum retractum</i> , <i>Calathu singratus</i> , <i>Platynus decentis</i> , <i>Pterostichus adstrictus</i> , <i>P. coracinus</i> , <i>P. pennsylvanicus</i> , <i>Sphaeroderus nitidicollis</i> , <i>Synuchus impunctatus</i>	Work et al. 2011
Coleoptera	Ground beetles	#	Bombi et al. 2019
Coleoptera	Saproxylc beetles	#	Gossner et al. 2016, Seibold et al. 2017
Coleoptera	Beetles	#	Müller et al. 2009 (b)
Hemiptera	Wēkiu bug	<i>Nysius wekiuicola</i>	Stephenson et al. 2017
Lepidoptera	Several species	<i>Coenonympha tullia</i> , <i>Celastrina ladon</i> , <i>Cupido amyntula</i> , <i>Vanessa cardui</i>	Hess et al. 2013
Lepidoptera	Several species	#	Aguirre-Gutierrez et al. 2017
Araneae	Spiders	#	Vierling et al. 2011
Decapoda	Red swamp crayfish	<i>Procambarus clarkii</i>	Paulson & Martin, 2014
Ostreida	Eastern oyster	<i>Crassostrea virginica</i>	Hogan & Reidenbach, 2019
Several	Several species	N/A	Kerfoot et al. 2019
Several	Arthropods	#	Müller et al. 2018, Müller et al. 2014
Several	Snails	#	Zellweger et al. 2015
<b>Reptiles</b>			
Crocodylia	Saltwater crocodile	<i>Crocodylus porosus</i>	Evans et al. 2017
Squamata	Several species	<i>Lacerta schreiberi</i> , <i>Timon lepidus</i> , <i>Podarcis bocagei</i> , <i>P. hispanica</i>	Sillero & Goncalves-Seco, 2014
Testudines	Kemp's ridley sea turtle	<i>Lepidochelys kempii</i>	Culver et al. 2020
Testudines	Greek tortoise	<i>Testudo graeca</i>	Bernheim et al. 2019
<b>Amphibians</b>			

Several	Several species	N/A	Julian et al. 2009, Wu et al. 2014, Riley et al. 2017
<b>Bryophytes, lichens &amp; fungi</b>			
Several	Bryophytes	<i>Sphagnum</i> spp.	Korpela et al. 2020
Several	Bryophytes	<i>Campylophyllum hispidulum</i> , <i>Chiloscyphus pallescens</i> , <i>Lophocolea heterophylla</i> , <i>Plagiomnium drummondii</i> , <i>Scapani aglaucocephala</i> , <i>Pleurozium schreberi</i> , <i>Hylocomium splendens</i> , <i>Aulacomnium palustre</i> , <i>Thuidium recognitum</i> , <i>Brachythecium campestre</i> , <i>Orthotrichum speciosum</i> , <i>Pylaisia polyantha</i> , <i>Ptilidium pulcherrimum</i> , <i>Dicranum fragilifolium</i>	Bartels et al. 2018
Several	Bryophytes	<i>Sphagnum</i> spp., <i>Pleurozium</i> spp., <i>Hylocomium</i> spp.	Peckham et al. 2009
Several	Bryophytes	<i>Pleurozium schreberi</i> , <i>Hylocomium splendens</i> , <i>Sphagnum girgensohnii</i> , <i>Polytrichum commune</i>	Kristensen et al. 2015
Several	Lichens	<i>Cladonia</i> & <i>Cetraria</i> spp	Keim et al. 2017
Several	Lichens (Biological soil crust)	N/A	Rodriguez-Caballero et al. 2019
Several	Fungi	#	Peura et al. 2016, Thers et al. 2017
Several	Several species	N/A	Langford et al. 2016, Moeslund et al. 2019
<b>Several taxonomic groups</b>			
Birds, bats, mammals, amphibians		Cooper et al. 2020	
Birds, bats, invertebrates, bryophytes, lichens, fungi		Bae et al. 2019	
Birds, mammals (bats)		Renner et al. 2018	
Birds, invertebrates (butterflies)		Zellweger et al. 2017	
Birds, invertebrates (insects)		Zellweger et al. 2016	
Birds, invertebrates (beetles)		Lindberg et al. 2015	
Birds, mammals, amphibians		Roll et al. 2015	

**Annex 2:** List of the reviewed articles included in the systematic review (173).

Note that all these articles have not been cited in the main text.

1. Adams, B.T. & Matthews, S.N. (2019). Diverse temperate forest bird assemblages demonstrate closer correspondence to plant species composition than vegetation structure. *Ecography*, 42(10): 1752-1764. doi: 10.1111/ecog.04487.
2. Aguirre-Gutierrez, J., Wallis DeVries, M.F., Marshall, L., van't Zelfde, M., Villalobos-Arambula, A.R., Boekelo, B., Bartholomeus, H., Franzen, M. & Biesmeijer, J.C. (2017). Butterflies show different functional and species diversity in relationship to vegetation structure and land use. *Global Ecology and Biogeography*, 26(10): 1126-1137. doi: 10.1111/geb.12622.
3. Amaral, K.E., Palace, M., O'Brien, K.M., Fenderson, L.E. & Kovach, A.I. (2016). Anthropogenic Habitats Facilitate Dispersal of an Early Successional Obligate: Implications for Restoration of an Endangered Ecosystem. *Plos One*, 11(3): e0148842. doi: 10.1371/journal.pone.0148842.
4. Asner, G.P., Owen-Smith, N., Loarie, S.R., Davies, A.B., Le Roux, E. & Levick, S.R. (2015). Habitat differences do not explain population declines of sable antelope in an African savanna. *Journal of Zoology*, 297(3): 225-234. doi: 10.1111/jzo.12269.
5. Bae, S., Reineking, B., Ewald, M. & Mueller, J. (2014). Comparison of airborne lidar, aerial photography, and field surveys to model the habitat suitability of a cryptic forest species - the hazel grouse. *International Journal of Remote Sensing*, 35(17): 6469-6489. doi: 10.1080/01431161.2014.955145.
6. Bae, S., Mueller, J., Lee, D., Vierling, K.T., Vogeler, J.C., Vierling, L.A., Hudak, A.T., Latifi, H. & Thorn, S. (2018). Taxonomic, functional, and phylogenetic diversity of bird assemblages are oppositely associated to productivity and heterogeneity in temperate forests. *Remote Sensing of Environment*, 215: 145-156. doi: 10.1016/j.rse.2018.05.031.
7. Bae, S., Levick, S.R., Heidrich, L., Magdon, P., Leutner, B.F., Wöllauer, S., Serebryanyk, A., Nauss, T., Krzystek, P., Gossner, M.M., Schall, P., Heibl, C., Bässler, C., Doerfler, I., Schulze, E., Krah, F., Culmsee, H., Jung, K., Heurich, M., Fischer, M., Seibold, S., Thorn, S., Gerlach, T., Hothorn, T., Weisser, W.W. & Müller, J. (2019). Radar vision in the mapping of forest biodiversity from space. *Nature Communications*, 10(1): 4757. doi: 10.1038/s41467-019-12737-x.
8. Barnes, K.W., Islam, K. & Auer, S.A. (2016). Integrating LIDAR-derived canopy structure into cerulean warbler habitat models. *Journal of Wildlife Management*, 80(1): 101-116. doi: 10.1002/jwmg.995.
9. Bartels, S.F., Caners, R.T., Ogilvie, J., White, B. & Macdonald, S.E. (2018). Relating Bryophyte Assemblages to a Remotely Sensed Depth-to-Water Index in Boreal Forests. *Frontiers in Plant Science*, 9: 858. doi: 10.3389/fpls.2018.00858.
10. Bellamy, P.E., Hill, R.A., Rothery, P., Hinsley, S.A., Fuller, R.J. & Broughton, R.K. (2009). Willow Warbler *Phylloscopus trochilus* habitat in woods with different structure and management in southern England. *Bird Study*, 56: 338-348. doi: 10.1080/00063650902806914.
11. Bernheim, M., Shanas, U. & Avi Bar-Massada (2019). Soil Type Impacts Macrohabitat Selection and Spatiotemporal Activity Patterns of *Testudo graeca* in an

- Eastern Mediterranean Ecosystem. *Copeia*, 107(2): 332-340. doi: 10.1643/CE-18-066R1.
12. Blomdahl, E.M., Thompson, C.M., Kane, J.R., Kane, V.R., Churchill, D., Moskal, L.M. & Lutz, J.A. (2019). Forest structure predictive of fisher (*Pekania pennanti*) dens exists in recently burned forest in Yosemite, California, USA. *Forest Ecology and Management*, 444: 174-186. doi: 10.1016/j.foreco.2019.04.024.
  13. Boelman, N.T., Holbrook, J.D., Greaves, H.E., Krause, J.S., Chmura, H.E., Magney, T.S., Perez, J.H., Eitel, J.U.H., Gough, L., Vierling, K.T., Wingfield, J.C. & Vierling, L.A. (2016). Airborne laser scanning and spectral remote sensing give a bird's eye perspective on arctic tundra breeding habitat at multiple spatial scales. *Remote Sensing of Environment*, 184: 337-349. doi: 10.1016/j.rse.2016.07.012.
  14. Boelman, N.T., Asner, G.P., Hart, P.J. & Martin, R.E. (2007). Multi-trophic invasion resistance in Hawaii: Bioacoustics, field surveys, and airborne remote sensing. *Ecological Applications*, 17(8): 2137-2144. doi: 10.1890/07-0004.1.
  15. Bombi, P., Gnetti, V., D'Andrea, E., De Cinti, B., Taglianti, A.V., Bologna, M.A. & Matteucci, G. (2019). Identifying priority sites for insect conservation in forest ecosystems at high resolution: the potential of LiDAR data. *Journal of Insect Conservation*, 23(4): 689-698. doi: 10.1007/s10841-019-00162-w.
  16. Bradbury, R.B., Hill, R.A., Mason, D.C., Hinsley, S.A., Wilson, J.D., Balzter, H., Anderson, G.Q.A., Whittingham, M.J., Davenport, I.J. & Bellamy, P.E. (2005). Modelling relationships between birds and vegetation structure using airborne LiDAR data: a review with case studies from agricultural and woodland environments. *Ibis*, 147(3): 443-452. doi: 10.1111/j.1474-919x.2005.00438.x.
  17. Broughton, R.K., Hinsley, S.A., Bellamy, P.E., Hill, R.A. & Rothery, P. (2006). Marsh Tit *Poecile palustris* territories in a British broad-leaved wood. *Ibis*, 148(4): 744-752. doi: 10.1111/j.1474-919X.2006.00583.x.
  18. Broughton, R.K., Hill, R.A., Freeman, S.N., Bellamy, P.E. & Hinsley, S.A. (2012a). Describing Habitat Occupation by Woodland Birds with Territory Mapping and Remotely Sensed Data: An Example using the Marsh Tit (*Poecile palustris*). *The Condor*, 114(4): 812-822. doi: 10.1525/cond.2012.110171.
  19. Broughton, R.K., Hill, R.A., Henderson, L.J., Bellamy, P.E. & Hinsley, S.A. (2012b). Patterns of nest placement in a population of Marsh Tits *Poecile palustris*. *Journal of Ornithology*, 153(3): 735-746. doi: 10.1007/s10336-011-0790-2.
  20. Broughton, R.K., Bellamy, P.E., Hill, R.A. & Hinsley, S.A. (2014). Winter habitat selection by Marsh Tits *Poecile palustris* in a British woodland. *Bird Study*, 61(3): 404-412. doi: 10.1080/00063657.2014.934193.
  21. Carr, A., Zeale, M.R.K., Weatherall, A., Froidevaux, J.S.P. & Jones, G. (2018). Ground-based and LiDAR-derived measurements reveal scale-dependent selection of roost characteristics by the rare tree-dwelling bat *Barbastella barbastellus*. *Forest Ecology and Management*, 417: 237-246. doi: 10.1016/j.foreco.2018.02.041.
  22. Carrasco, L., Giam, X., Papes, M. & Sheldon, K.S. (2019). Metrics of Lidar-Derived 3D Vegetation Structure Reveal Contrasting Effects of Horizontal and Vertical Forest Heterogeneity on Bird Species Richness. *Remote Sensing*, 11(7): 743. doi: 10.3390/rs11070743.
  23. Casas, A., Garcia, M., Siegel, R.B., Koltunov, A., Ramirez, C. & Ustin, S. (2016). Burned forest characterization at single-tree level with airborne laser scanning for

- assessing wildlife habitat. *Remote Sensing of Environment*, 175: 231-241. doi: 10.1016/j.rse.2015.12.044.
24. Clawges, R., Vierling, K., Vierling, L. & Rowell, E. (2008). The use of airborne lidar to assess avian species diversity, density, and occurrence in a pine/aspen forest. *Remote Sensing of Environment*, 112(5): 2064-2073. doi: 10.1016/j.rse.2007.08.023.
  25. Cooper, W.J., McShea, W.J., Luther, D.A. & Forrester, T. (2020). Incorporating local habitat heterogeneity and productivity measures when modelling vertebrate richness. *Environmental Conservation*, 47(1): 7-14. doi: 10.1017/S0376892919000328.
  26. Coops, N.C., Duffe, J. & Koot, C. (2010). Assessing the utility of lidar remote sensing technology to identify mule deer winter habitat. *Canadian Journal of Remote Sensing*, 36(2): 81-88. doi: 10.5589/m10-029.
  27. Coops, N.C., Tompaski, P., Nijland, W., Rickbeil, G.J.M., Nielsen, S.E., Bater, C.W. & Stadt, J.J. (2016). A forest structure habitat index based on airborne laser scanning data. *Ecological Indicators*, 67: 346-357. doi: 10.1016/j.ecolind.2016.02.057.
  28. Culver, M., Gibeaut, J.C., Shaver, D.J., Tissot, P. & Starek, M. (2020). Using Lidar Data to Assess the Relationship Between Beach Geomorphology and Kemp's Ridley (*Lepidochelys kempii*) Nest Site Selection Along Padre Island, TX, United States. *Frontiers in Marine Science*, 7: 214.
  29. Davenport, I.J., Bradbury, R.B., Anderson, G., Hayman, G., Krebs, J.R., Mason, D.C., Wilson, J.D. & Veck, N.J. (2000). Improving bird population models using airborne remote sensing. *International Journal of Remote Sensing*, 21(13-14): 2705-2717. doi: 10.1080/01431160050110241.
  30. Davies, A.B., Levick, S.R., Asner, G.P., Robertson, M.P., van Rensburg, B.J. & Parr, C.L. (2014). Spatial variability and abiotic determinants of termite mounds throughout a savanna catchment. *Ecography*, 37(9): 852-862. doi: 10.1111/ecog.00532.
  31. Davies, A.B., Baldeck, C.A. & Asner, G.P. (2016). Termite mounds alter the spatial distribution of African savanna tree species. *Journal of Biogeography*, 43(2): 301-313. doi: 10.1111/jbi.12633.
  32. Davies, A.B., Tambling, C.J., Kerley, G.I.H. & Asner, G.P. (2016a). Effects of Vegetation Structure on the Location of Lion Kill Sites in African Thicket. *Plos One*, 11(2): e0149098. doi: 10.1371/journal.pone.0149098.
  33. Davies, A.B., Tambling, C.J., Kerley, G.I.H. & Asner, G.P. (2016b). Limited spatial response to direct predation risk by African herbivores following predator reintroduction. *Ecology and Evolution*, 6(16): 5728-5748. doi: 10.1002/ece3.2312.
  34. Davies, A.B., Oram, F., Ancrenaz, M. & Asner, G.P. (2019). Combining behavioural and LiDAR data to reveal relationships between canopy structure and orangutan nest site selection in disturbed forests. *Biological Conservation*, 232: 97-107. doi: 10.1016/j.biocon.2019.01.032.
  35. de Thoisy, B., Fayad, I., Clement, L., Barrioz, S., Poirier, E. & Gond, V. (2016). Predators, Prey and Habitat Structure: Can Key Conservation Areas and Early Signs of Population Collapse Be Detected in Neotropical Forests?. *Plos One*, 11(11): e0165362. doi: 10.1371/journal.pone.0165362.
  36. Ehmke, G., Maguire, G.S., Bird, T., Ierodiaconou, D. & Weston, M.A. (2016). An obligate beach bird selects sub-, inter- and supra-tidal habitat elements. *Estuarine Coastal and Shelf Science*, 181: 266-276. doi: 10.1016/j.ecss.2016.08.050.
  37. Eldegard, K., Dirksen, J.W., Orka, H.O., Halvorsen, R., Naesset, E., Gobakken, T. & Ohlson, M. (2014). Modelling bird richness and bird species presence in a boreal

- forest reserve using airborne laser-scanning and aerial images. *Bird Study*, 61(2): 204-219. doi: 10.1080/00063657.2014.885492.
38. Evans, L.J., Davies, A.B., Goossens, B. & Asner, G.P. (2017). Riparian vegetation structure and the hunting behavior of adult estuarine crocodiles. *Plos One*, 12(10): e0184804. doi: 10.1371/journal.pone.0184804.
  39. Ewald, M., Dupke, C., Heurich, M., Mueller, J. & Reineking, B. (2014). LiDAR Remote Sensing of Forest Structure and GPS Telemetry Data Provide Insights on Winter Habitat Selection of European Roe Deer. *Forests*, 5(6): 1374-1390. doi: 10.3390/f5061374.
  40. Ficetola, G.F., Bonardi, A., Mucher, C.A., Gilissen, N.L.M. & Padoa-Schioppa, E. (2014). How many predictors in species distribution models at the landscape scale? Land use versus LiDAR-derived canopy height. *International Journal of Geographical Information Science*, 28(8): 1723-1739. doi: 10.1080/13658816.2014.891222.
  41. Finnegan, L., Pigeon, K.E., Cranston, J., Hebblewhite, M., Musiani, M., Neufeld, L., Schmiegelow, F., Duval, J. & Stenhouse, G.B. (2018). Natural regeneration on seismic lines influences movement behaviour of wolves and grizzly bears. *Plos One*, 13(4): e0195480. doi: 10.1371/journal.pone.0195480.
  42. Flaherty, S., Lurz, P.W.W. & Patenaude, G. (2014). Use of LiDAR in the conservation management of the endangered red squirrel (*Sciurus vulgaris* L.). *Journal of Applied Remote Sensing*, 8: 083592. doi: 10.1117/1.JRS.8.083592.
  43. Flaspohler, D.J., Giardina, C.P., Asner, G.P., Hart, P., Price, J., Lyons, C.K. & Castaneda, X. (2010). Long-term effects of fragmentation and fragment properties on bird species richness in Hawaiian forests. *Biological Conservation*, 143(2): 280-288. doi: 10.1016/j.biocon.2009.10.009.
  44. Fotis, A.T., Patel, S. & Chavez, A.S. (2020). Habitat-based isolating barriers are not strong in the speciation of ecologically divergent squirrels (*Tamiasciurus douglasii* and *T. hudsonicus*). *Behavioral Ecology and Sociobiology*, 74(3): 32. doi: 10.1007/s00265-020-2814-5.
  45. Froidevaux, J.S.P., Zellweger, F., Bollmann, K., Jones, G. & Obrist, M.K. (2016). From field surveys to LiDAR: Shining a light on how bats respond to forest structure. *Remote Sensing of Environment*, 175: 242-250. doi: 10.1016/j.rse.2015.12.038.
  46. Gallagher, C.V., Keane, J.J., Shaklee, P.A., Kramer, H.A. & Gerrard, R. (2019). Spotted owl foraging patterns following fuels treatments, Sierra Nevada, California. *Journal of Wildlife Management*, 83(2): 487-501. doi: 10.1002/jwmg.21586.
  47. Garabedian, J.E., McGaughey, R.J., Reutebuch, S.E., Parresol, B.R., Kilgo, J.C., Moorman, C.E. & Peterson, M.N. (2014). Quantitative analysis of woodpecker habitat using high-resolution airborne LiDAR estimates of forest structure and composition. *Remote Sensing of Environment*, 145: 68-80. doi: 10.1016/j.rse.2014.01.022.
  48. Garabedian, J.E., Moorman, C.E., Peterson, M.N. & Kilgo, J.C. (2017). Use of LiDAR to define habitat thresholds for forest bird conservation. *Forest Ecology and Management*, 399: 24-36. doi: 10.1016/j.foreco.2017.05.024.
  49. Garabedian, J.E., Moorman, C.E., Peterson, M.N. & Kilgo, J.C. (2018). Relative importance of social factors, conspecific density, and forest structure on space use by the endangered Red-cockaded Woodpecker: A new consideration for habitat restoration. *Condor*, 120(2): 305-318. doi: 10.1650/CONDOR-17-211.1.

50. Garabedian, J.E., Moorman, C.E., Peterson, M.N. & Kilgo, J.C. (2020). Effects of group size and group density on trade-offs in resource selection by a group-territorial central-place foraging woodpecker. *Ibis*, 162(2): 477-491. doi: 10.1111/ibi.12733.
51. Garcia-Feced, C., Tempel, D.J. & Kelly, M. (2011). LiDAR as a Tool to Characterize Wildlife Habitat: California Spotted Owl Nesting Habitat as an Example. *Journal of Forestry*, 109(8): 436-443.
52. Glad, A., Reineking, B., Montadert, M., Depraz, A. & Monnet, J. (2020). Assessing the performance of object-oriented LiDAR predictors for forest bird habitat suitability modeling. *Remote Sensing in Ecology and Conservation*, 6(1): 5-19. doi: 10.1002/rse2.117.
53. Goetz, S., Steinberg, D., Dubayah, R. & Blair, B. (2007). Laser remote sensing of canopy habitat heterogeneity as a predictor of bird species richness in an eastern temperate forest, USA. *Remote Sensing of Environment*, 108(3): 254-263. doi: 10.1016/j.rse.2006.11.016.
54. Goetz, S.J., Steinberg, D., Betts, M.G., Holmes, R.T., Doran, P.J., Dubayah, R. & Hofton, M. (2010). Lidar remote sensing variables predict breeding habitat of a Neotropical migrant bird. *Ecology*, 91(6): 1569-1576. doi: 10.1890/09-1670.1.
55. Goetz, S.J., Sun, M., Zolkos, S., Hansen, A. & Dubayah, R. (2014). The relative importance of climate and vegetation properties on patterns of North American breeding bird species richness. *Environmental Research Letters*, 9(3): 034013. doi: 10.1088/1748-9326/9/3/034013.
56. Goodwin, C.E.D., Hodgson, D.J., Bailey, S., Bennie, J. & McDonald, R.A. (2018). Habitat preferences of hazel dormice *Muscardinus avellanarius* and the effects of tree-felling on their movement. *Forest Ecology and Management*, 427: 190-199. doi: 10.1016/j.foreco.2018.03.035.
57. Gossner, M.M., Wende, B., Levick, S., Schall, P., Floren, A., Linsenmair, K.E., Steffan-Dewenter, I., Schulze, E. & Weisser, W.W. (2016). Deadwood enrichment in European forests - Which tree species should be used to promote saproxylic beetle diversity?. *Biological Conservation*, 201: 92-102. doi: 10.1016/j.biocon.2016.06.032.
58. Graf, R.F., Mathys, L. & Bollmann, K. (2009). Habitat assessment for forest dwelling species using LiDAR remote sensing: Capercaillie in the Alps. *Forest Ecology and Management*, 257(1): 160-167. doi: 10.1016/j.foreco.2008.08.021.
59. Hagar, J.C., Eskelson, B.N.I., Haggerty, P.K., Nelson, S.K. & Vesely, D.G. (2014). Modeling Marbled Murrelet (*Brachyramphus marmoratus*) Habitat Using LiDAR-Derived Canopy Data. *Wildlife Society Bulletin*, 38(2): 237-249. doi: 10.1002/wsb.407.
60. Hagar, J.C., Yost, A. & Haggerty, P.K. (2020). Incorporating LiDAR metrics into a structure-based habitat model for a canopy-dwelling species. *Remote Sensing of Environment*, 236: 111499. doi: 10.1016/j.rse.2019.111499.
61. Hatfield, J.S., Reynolds, M.H., Seavy, N.E. & Krause, C.M. (2012). Population Dynamics of Hawaiian Seabird Colonies Vulnerable to Sea-Level Rise. *Conservation Biology*, 26(4): 667-678. doi: 10.1111/j.1523-1739.2012.01853.x.
62. Hatten, J.R. (2014). Mapping and monitoring Mount Graham red squirrel habitat with Lidar and Landsat imagery. *Ecological Modelling*, 289: 106-123. doi: 10.1016/j.ecolmodel.2014.07.004.
63. Hess, A.N., Falkowski, M.J., Webster, C.R., Storer, A.J., Pocewicz, A. & Martinuzzi, S. (2013). Employing lidar data to identify butterfly habitat characteristics of four

- contrasting butterfly species across a diverse landscape. *Remote Sensing Letters*, 4(4): 354-363. doi: 10.1080/2150704X.2012.734932.
64. Hill, R.A. & Hinsley, S.A. (2015). Airborne Lidar for Woodland Habitat Quality Monitoring: Exploring the Significance of Lidar Data Characteristics when Modelling Organism-Habitat Relationships. *Remote Sensing*, 7(4): 3446-3466. doi: 10.3390/rs70403446.
65. Hinsley, S.A., Hill, R.A., Gaveau, D.L.A. & Bellamy, P.E. (2002). Quantifying woodland structure and habitat quality for birds using airborne laser scanning. *Functional Ecology*, 16(6): 851-857. doi: 10.1046/j.1365-2435.2002.00697.x.
66. Hinsley, S.A., Hill, R.A., Bellamy, P.E. & Balzter, H. (2006). The application of lidar in woodland bird ecology: Climate, canopy structure, and habitat quality. *Photogrammetric Engineering and Remote Sensing*, 72(12): 1399-1406. doi: 10.14358/PERS.72.12.1399.
67. Hinsley, S.A., Hill, R.A., Bellamy, P.E., Harrison, N.M., Speakman, J.R., Wilson, A.K. & Ferns, P.N. (2008). Effects of structural and functional habitat gaps on breeding woodland birds: working harder for less. *Landscape Ecology*, 23(5): 615-626. doi: 10.1007/s10980-008-9225-8.
68. Hinsley, S.A., Hill, R.A., Bellamy, P., Broughton, R.K., Harrison, N.M., Mackenzie, J.A., Speakman, J.R. & Ferns, P.N. (2009a). Do Highly Modified Landscapes Favour Generalists at the Expense of Specialists? An Example using Woodland Birds. *Landscape Research*, 34(5): 509-526. doi: 10.1080/01426390903177276.
69. Hinsley, S.A., Hill, R.A., Fuller, R.J., Bellamy, P.E. & Rothery, P. (2009b). Bird species distributions across woodland canopy structure gradients. *Community Ecology*, 10(1): 99-110. doi: 10.1556/ComEc.10.2009.1.12.
70. Hogan, S. & Reidenbach, M.A. (2019). Quantifying and mapping intertidal oyster reefs utilizing LiDAR-based remote sensing. *Mar Ecol Prog Ser*, 630: 83-99.
71. Huber, N., Kienast, F., Ginzler, C. & Pasinelli, G. (2016). Using remote-sensing data to assess habitat selection of a declining passerine at two spatial scales. *Landscape Ecology*, 31(9): 1919-1937. doi: 10.1007/s10980-016-0370-1.
72. Huber, N., Kery, M. & Pasinelli, G. (2017). Occupancy dynamics of the Wood Warbler *Phylloscopus sibilatrix* assessed with habitat and remote sensing data. *Ibis*, 159(3): 623-637. doi: 10.1111/ibi.12472.
73. Hull, I.T. & Shipley, L.A. (2019). Testing the Ability of Airborne LiDAR to Measure Forage Resources for Wild Ungulates in Conifer Forests. *Journal of Forestry*, 117(5): 492-503. doi: 10.1093/jofore/fvz040.
74. Jaime-Gonzalez, C., Acebes, P., Mateos, A. & Mezquida, E.T. (2017). Bridging gaps: On the performance of airborne LiDAR to model wood mouse-habitat structure relationships in pine forests. *Plos One*, 12(8): e0182451. doi: 10.1371/journal.pone.0182451.
75. Jenkins, J.M.A., Lesmeister, D.B., Wiens, J.D., Kane, J.T., Kane, V.R. & Verschuyt, J. (2019). Three-dimensional partitioning of resources by congeneric forest predators with recent sympatry. *Scientific Reports*, 9: 6036. doi: 10.1038/s41598-019-42426-0.
76. Jirinec, V., Isdell, R.E. & Leu, M. (2016). Prey availability and habitat structure explain breeding space use of a migratory songbird. *Condor*, 118(2): 309-328. doi: 10.1650/CONDOR-15-140.1.
77. Jones, T.G., Arcese, P., Sharma, T. & Coops, N.C. (2013). Describing avifaunal richness with functional and structural bioindicators derived from advanced

- airborne remotely sensed data. *International Journal of Remote Sensing*, 34(8): 2689-2713. doi: 10.1080/01431161.2012.750019.
78. Julian, J.T., Young, J.A., Jones, J.W., Snyder, C.D. & Wright, C.W. (2009). The use of local indicators of spatial association to improve LiDAR-derived predictions of potential amphibian breeding ponds. *Journal of Geographical Systems*, 11(1): 89-106. doi: 10.1007/s10109-008-0074-4.
79. Jung, K., Kaiser, S., Boehm, S., Nieschulze, J. & Kalko, E.K.V. (2012). Moving in three dimensions: effects of structural complexity on occurrence and activity of insectivorous bats in managed forest stands. *Journal of Applied Ecology*, 49(2): 523-531. doi: 10.1111/j.1365-2664.2012.02116.x.
80. Kawaguchi, T. & Desrochers, A. (2017). Short-term changes in spatial distribution pattern of an herbivore in response to accumulating snow. *Canadian Journal of Zoology*, 95(12): 949-954. doi: 10.1139/cjz-2016-0287.
81. Keim, J.L., DeWitt, P.D., Fitzpatrick, J.J. & Jenni, N.S. (2017). Estimating plant abundance using inflated beta distributions: Applied learnings from a lichen-caribou ecosystem. *Ecology and Evolution*, 7(2): 486-493. doi: 10.1002/ece3.2625.
82. Kellner, K.F., Ruhl, P.J., Dunning, J.B., Jr., Barnes, K.W., Saunders, M.R. & Swihart, R.K. (2018). Local-scale Habitat Components Driving Bird Abundance in Eastern Deciduous Forests. *American Midland Naturalist*, 180(1): 52-65. doi: 10.1674/0003-0031-180.1.52.
83. Kerfoot, W.C., Hobmeier, M.M., Regis, R., Raman, V.K., Brooks, C.N., Shuchman, R., Sayers, M., Yousef, F. & Reif, M. (2019). Lidar (light detection and ranging) and benthic invertebrate investigations: Migrating tailings threaten Buffalo Reef in Lake Superior. *Journal of Great Lakes Research*, 45(5): 872-887. doi: 10.1016/j.jglr.2019.07.009.
84. Khanaposhtani, M.G., Gasc, A., Francomano, D., Villanueva-Rivera, L.J., Jung, J., Mossman, M.J. & Pijanowski, B.C. (2019). Effects of highways on bird distribution and soundscape diversity around Aldo Leopold's shack in Baraboo, Wisconsin, USA. *Landscape and Urban Planning*, 192: UNSP 103666. doi: 10.1016/j.landurbplan.2019.103666.
85. Klein, J., Haverkamp, P.J., Lindberg, E., Griesser, M. & Eggers, S. (2020). Remotely sensed forest understory density and nest predator occurrence interact to predict suitable breeding habitat and the occurrence of a resident boreal bird species. *Ecology and Evolution*, 10(4): 2238-2252. doi: 10.1002/ece3.6062.
86. Korpela, I., Haapanen, R., Korrensalo, A., Tuittila, E. & Vesala, T. (2020). Fine-resolution mapping of microforms of a boreal bog using aerial images and waveform-recording LiDAR. *Mires and Peat*, 26 doi: 10.19189/MaP.2018.OMB.388.
87. Kortmann, M., Hurst, J., Brinkmann, R., Heurich, M., Gonzalez, R.S., Mueller, J. & Thorn, S. (2018b). Beauty and the beast: how a bat utilizes forests shaped by outbreaks of an insect pest. *Animal Conservation*, 21(1): 21-30. doi: 10.1111/acv.12359.
88. Kortmann, M., Heurich, M., Latifi, H., Roesner, S., Seidl, R., Mueller, J. & Thorn, S. (2018a). Forest structure following natural disturbances and early succession provides habitat for two avian flagship species, capercaillie (*Tetrao urogallus*) and hazel grouse (*Tetrastes bonasia*). *Biological Conservation*, 226: 81-91. doi: 10.1016/j.biocon.2018.07.014.

89. Kristensen, T., Naesset, E., Ohlson, M., Bolstad, P.V. & Kolka, R. (2015). Mapping Above- and Below-Ground Carbon Pools in Boreal Forests: The Case for Airborne Lidar. *Plos One*, 10(10): e0138450. doi: 10.1371/journal.pone.0138450.
90. Kroeger, A.J., Moorman, C.E., Lashley, M.A., Chitwood, M.C., Harper, C.A. & DePerno, C.S. (2020). White-tailed deer use of overstory hardwoods in longleaf pine woodlands. *Forest Ecology and Management*, 464: 118046. doi: 10.1016/j.foreco.2020.118046.
91. Lacki, M.J., Dodd, L.E., Skowronski, N.S., Dickinson, M.B. & Rieske, L.K. (2017). Relationships among burn severity, forest canopy structure and bat activity from spring burns in oak-hickory forests. *International Journal of Wildland Fire*, 26(11): 963-972. doi: 10.1071/WF16159.
92. Langford, Z., Kumar, J., Hoffman, F.M., Norby, R.J., Wulschleger, S.D., Sloan, V.L. & Iversen, C.M. (2016). Mapping Arctic Plant Functional Type Distributions in the Barrow Environmental Observatory Using WorldView-2 and LiDAR Datasets. *Remote Sensing*, 8(9): 733. doi: 10.3390/rs8090733.
93. Leivits, M. & Leivits, A. (2009). Use of sequential aerial photography and LiDAR for mapping Scots Pine (*Pinus sylvestris*) encroachment and change detection in bird habitats from 1950 to 2007 in Nigula Bog (Estonia). doi: 10.13140/RG.2.1.2668.8728.
94. Lesak, A.A., Radloff, V.C., Hawbaker, T.J., Pidgeon, A.M., Gobakken, T. & Contrucci, K. (2011). Modeling forest songbird species richness using LiDAR-derived measures of forest structure. *Remote Sensing of Environment*, 115(11): 2823-2835. doi: 10.1016/j.rse.2011.01.025.
95. Levick, S.R., Asner, G.P., Chadwick, O.A., Khomo, L.M., Rogers, K.H., Hartshorn, A.S., Kennedy-Bowdoin, T. & Knapp, D.E. (2010). Regional insight into savanna hydrogeomorphology from termite mounds. *Nature Communications*, 1: 65. doi: 10.1038/ncomms1066.
96. Lindberg, E., Roberge, J., Johansson, T. & Hjalten, J. (2015). Can Airborne Laser Scanning (ALS) and Forest Estimates Derived from Satellite Images Be Used to Predict Abundance and Species Richness of Birds and Beetles in Boreal Forest?. *Remote Sensing*, 7(4): 4233-4252. doi: 10.3390/rs70404233.
97. Linnell, M.A., Davis, R.J., Lesmeister, D.B. & Swingle, J.K. (2017). Conservation and relative habitat suitability for an arboreal mammal associated with old forest. *Forest Ecology and Management*, 402: 1-11. doi: 10.1016/j.foreco.2017.07.004.
98. Loarie, S.R., Tambling, C.J. & Asner, G.P. (2013). Lion hunting behaviour and vegetation structure in an African savanna. *Animal Behaviour*, 85(5): 899-906. doi: 10.1016/j.anbehav.2013.01.018.
99. Lone, K., Loe, L.E., Gobakken, T., Linnell, J.D.C., Odden, J., Remmen, J. & Mysterud, A. (2014). Living and dying in a multi-predator landscape of fear: roe deer are squeezed by contrasting pattern of predation risk imposed by lynx and humans. *Oikos*, 123(6): 641-651. doi: 10.1111/j.1600-0706.2013.00938.x.
100. Lorenz, T.J., Vierling, K.T., Vogeler, J., Lonneker, J. & Aycrigg, J. (2015). Incorporating Shrub and Snag Specific LiDAR Data into GAP Wildlife Models. *Journal of Fish and Wildlife Management*, 6(2): 437-447. doi: 10.3996/092013-JFWM-064.
101. Mangan, A.O., Chestnut, T., Vogeler, J.C., Breckheimer, I.K., King, W.M., Bagnall, K.E. & Dugger, K.M. (2019). Barred Owls reduce occupancy and breeding

- propensity of Northern Spotted Owl in a Washington old-growth forest. *Condor*, 121(3): UNSP duz031. doi: 10.1093/condor/duz031.
102. Martinuzzi, S., Vierling, L.A., Gould, W.A., Falkowski, M.J., Evans, J.S., Hudak, A.T. & Vierling, K.T. (2009). Mapping snags and understory shrubs for a LiDAR-based assessment of wildlife habitat suitability. *Remote Sensing of Environment*, 113(12): 2533-2546. doi: 10.1016/j.rse.2009.07.002.
  103. Mason, D.C., Anderson, G.Q.A., Bradbury, R.B., Cobby, D.M., Davenport, I.J., Vandepoll, M. & Wilson, J.D. (2003). Measurement of habitat predictor variables for organism-habitat models using remote sensing and image segmentation. *International Journal of Remote Sensing*, 24(12): 2515-2532. doi: 10.1080/014311602100100848.
  104. McLean, K.A., Trainor, A.M., Asner, G.P., Crofoot, M.C., Hopkins, M.E., Campbell, C.J., Martin, R.E., Knapp, D.E. & Jansen, P.A. (2016). Movement patterns of three arboreal primates in a Neotropical moist forest explained by LiDAR-estimated canopy structure. *Landscape Ecology*, 31(8): 1849-1862. doi: 10.1007/s10980-016-0367-9.
  105. Meiman, S., Civco, D., Holsinger, K. & Elphick, C.S. (2012). Comparing Habitat Models Using Ground-Based and Remote Sensing Data: Saltmarsh Sparrow Presence Versus Nesting. *Wetlands*, 32(4): 725-736. doi: 10.1007/s13157-012-0306-8.
  106. Melin, M., Packalen, P., Matala, J., Mehtatalo, L. & Pusenius, J. (2013). Assessing and modeling moose (*Alces alces*) habitats with airborne laser scanning data. *International Journal of Applied Earth Observation and Geoinformation*, 23: 389-396. doi: 10.1016/j.jag.2012.11.004.
  107. Melin, M., Matala, J., Mehtatalo, L., Pusenius, J. & Packalen, P. (2016). Ecological dimensions of airborne laser scanning - Analyzing the role of forest structure in moose habitat use within a year. *Remote Sensing of Environment*, 173: 238-247. doi: 10.1016/j.rse.2015.07.025.
  108. Melin, M., Hinsley, S.A., Broughton, R.K., Bellamy, P. & Hill, R.A. (2018). Living on the edge: utilising lidar data to assess the importance of vegetation structure for avian diversity in fragmented woodlands and their edges. *Landscape Ecology*, 33(6): 895-910. doi: 10.1007/s10980-018-0639-7.
  109. Melin, M., Matala, J., Mehtatalo, L., Tiilikainen, R., Tikkanen, O., Maltamo, M., Pusenius, J. & Packalen, P. (2014). Moose (*Alces alces*) reacts to high summer temperatures by utilizing thermal shelters in boreal forests - an analysis based on airborne laser scanning of the canopy structure at moose locations. *Global Change Biology*, 20(4): 1115-1125. doi: 10.1111/gcb.12405.
  110. Melin, M., Hill, R.A., Bellamy, P.E. & Hinsley, S.A. (2019). On Bird Species Diversity and Remote Sensing-Utilizing Lidar and Hyperspectral Data to Assess the Role of Vegetation Structure and Foliage Characteristics as Drivers of Avian Diversity. *Ieee Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 12(7): 2270-2278. doi: 10.1109/JSTARS.2019.2906940.
  111. Milanese, P., Holderegger, R., Bollmann, K., Gugerli, F. & Zellweger, F. (2017). Three-dimensional habitat structure and landscape genetics: a step forward in estimating functional connectivity. *Ecology*, 98(2): 393-402. doi: 10.1002/ecy.1645.
  112. Moeslund, J.E., Zlinszky, A., Ejrnaes, R., Brunbjerg, A.K., Bocher, P.K., Svenning, J. & Normand, S. (2019). Light detection and ranging explains diversity of plants, fungi, lichens, and bryophytes across multiple habitats and large

- geographic extent. *Ecological Applications*, 29(5): UNSP e01907. doi: 10.1002/eap.1907.
113. Mononen, L., Auvinen, A.-., Packalen, P., Virkkala, R., Valbuena, R., Bohlin, I., Valkama, J. & Vihervaara, P. (2018). Usability of citizen science observations together with airborne laser scanning data in determining the habitat preferences of forest birds. *Forest Ecology and Management*, 430: 498-508. doi: 10.1016/j.foreco.2018.08.040.
  114. Moreira-Arce, D., Vergara, P.M., Boutin, S., Carrasco, G., Briones, R., Soto, G.E. & Jimenez, J.E. (2016). Mesocarnivores respond to fine-grain habitat structure in a mosaic landscape comprised by commercial forest plantations in southern Chile. *Forest Ecology and Management*, 369: 135-143. doi: 10.1016/j.foreco.2016.03.024.
  115. Müller, J. & Brandl, R. (2009). Assessing biodiversity by remote sensing in mountainous terrain: the potential of LiDAR to predict forest beetle assemblages. *Journal of Applied Ecology*, 46(4): 897-905. doi: 10.1111/j.1365-2664.2009.01677.x.
  116. Müller, J., Moning, C., Baessler, C., Heurich, M. & Brandl, R. (2009). Using airborne laser scanning to model potential abundance and assemblages of forest passerines. *Basic and Applied Ecology*, 10(7): 671-681. doi: 10.1016/j.baae.2009.03.004.
  117. Müller, J., Stadler, J. & Brandl, R. (2010). Composition versus physiognomy of vegetation as predictors of bird assemblages: The role of lidar. *Remote Sensing of Environment*, 114(3): 490-495. doi: 10.1016/j.rse.2009.10.006.
  118. Müller, J., Brandl, R., Buchner, J., Pretzsch, H., Seifert, S., Straetz, C., Veith, M. & Fenton, B. (2013). From ground to above canopy-Bat activity in mature forests is driven by vegetation density and height. *Forest Ecology and Management*, 306: 179-184. doi: 10.1016/j.foreco.2013.06.043.
  119. Müller, J., Bae, S., Roeder, J., Chao, A. & Didham, R.K. (2014). Airborne LiDAR reveals context dependence in the effects of canopy architecture on arthropod diversity. *Forest Ecology and Management*, 312: 129-137. doi: 10.1016/j.foreco.2013.10.014.
  120. Müller, J., Brandl, R., Braendle, M., Foerster, B., de Araujo, B.C., Gossner, M.M., Ladas, A., Wagner, M., Maraun, M., Schall, P., Schmidt, S., Heurich, M., Thorn, S. & Seibold, S. (2018). LiDAR-derived canopy structure supports the more-individuals hypothesis for arthropod diversity in temperate forests. *Oikos*, 127(6): 814-824. doi: 10.1111/oik.04972.
  121. Nelson, R., Keller, C. & Ratnaswamy, M. (2005). Locating and estimating the extent of Delmarva fox squirrel habitat using an airborne LiDAR profiler. *Remote Sensing of Environment*, 96(3-4): 292-301. doi: 10.1016/j.rse.2005.02.012.
  122. North, M.P., Kane, J.T., Kane, V.R., Asner, G.P., Berigan, W., Churchill, D.J., Conway, S., Gutierrez, R.J., Jeronimo, S., Keane, J., Koltunov, A., Mark, T., Moskal, M., Munton, T., Peery, Z., Ramirez, C., Sollmann, R., White, A. & Whitmore, S. (2017). Cover of tall trees best predicts California spotted owl habitat. *Forest Ecology and Management*, 405: 166-178. doi: 10.1016/j.foreco.2017.09.019.
  123. Palminteri, S., Powell, G.V.N., Asner, G.P. & Peres, C.A. (2012). LiDAR measurements of canopy structure predict spatial distribution of a tropical mature forest primate. *Remote Sensing of Environment*, 127: 98-105. doi: 10.1016/j.rse.2012.08.014.
  124. Paulson, E.L. & Martin, A.P. (2014). Discerning invasion history in an ephemeral connected system: landscape genetics of *Procambarus clarkii* in Ash

- Meadows, Nevada. *Biological Invasions*, 16(8): 1719-1734. doi: 10.1007/s10530-013-0621-x.
125. Peckham, S.D., Ahl, D.E. & Gower, S.T. (2009). Bryophyte cover estimation in a boreal black spruce forest using airborne lidar and multispectral sensors. *Remote Sensing of Environment*, 113(6): 1127-1132. doi: 10.1016/j.rse.2009.02.008.
  126. Peura, M., Gonzalez, R.S., Mueller, J., Heurich, M., Vierling, L.A., Moenkkoenen, M. & Baessler, C. (2016). Mapping a 'cryptic kingdom': Performance of lidar derived environmental variables in modelling the occurrence of forest fungi. *Remote Sensing of Environment*, 186: 428-438. doi: 10.1016/j.rse.2016.09.003.
  127. Redhead, J.W., Pywell, R.F., Bellamy, P.E., Broughton, R.K., Hill, R.A. & Hinsley, S.A. (2013). Great tits *Parus major* and blue tits *Cyanistes caeruleus* as indicators of agri-environmental habitat quality. *Agriculture Ecosystems & Environment*, 178: 31-38. doi: 10.1016/j.agee.2013.06.015.
  128. Renner, S.C., Baur, S., Possler, A., Winkler, J., Kalko, E.K.V., Bates, P.J.J. & Mello, M.A.R. (2012). Food Preferences of Winter Bird Communities in Different Forest Types. *Plos One*, 7(12): e53121. doi: 10.1371/journal.pone.0053121.
  129. Renner, S.C., Luedtke, B., Kaiser, S., Kienle, J., Schaefer, H.M., Segelbacher, G., Tschapka, M. & Santiago-Alarcon, D. (2016). Forests of opportunities and mischief: disentangling the interactions between forests, parasites and immune responses. *International Journal for Parasitology*, 46(9): 571-579. doi: 10.1016/j.ijpara.2016.04.008.
  130. Renner, S.C., Suarez-Rubio, M., Kaiser, S., Nieschulze, J., Kalko, E.K.V., Tschapka, M. & Jung, K. (2018). Divergent response to forest structure of two mobile vertebrate groups. *Forest Ecology and Management*, 415: 129-138. doi: 10.1016/j.foreco.2018.02.028.
  131. Riley, J.W., Calhoun, D.L., Barichivich, W.J. & Walls, S.C. (2017). Identifying Small Depressional Wetlands and Using a Topographic Position Index to Infer Hydroperiod Regimes for Pond-Breeding Amphibians. *Wetlands*, 37(2): 325-338. doi: 10.1007/s13157-016-0872-2.
  132. Rodriguez-Caballero, E., Raul Roman, J., Chamizo, S., Roncero Ramos, B. & Canton, Y. (2019). Biocrust landscape-scale spatial distribution is strongly controlled by terrain attributes: Topographic thresholds for colonization in a semiarid badland system. *Earth Surface Processes and Landforms*, 44(14): 2771-2779. doi: 10.1002/esp.4706.
  133. Roll, U., Geffen, E. & Yom-Tov, Y. (2015). Linking vertebrate species richness to tree canopy height on a global scale. *Global Ecology and Biogeography*, 24(7): 814-825. doi: 10.1111/geb.12325.
  134. Sasaki, T., Imanishi, J., Fukui, W. & Morimoto, Y. (2016). Fine-scale characterization of bird habitat using airborne LiDAR in an urban park in Japan. *Urban Forestry & Urban Greening*, 17: 16-22. doi: 10.1016/j.ufug.2016.03.007.
  135. Schooler, S.L. & Zald, H.S.J. (2019). Lidar Prediction of Small Mammal Diversity in Wisconsin, USA. *Remote Sensing*, 11(19) doi: 10.3390/rs11192222.
  136. Seaman, D.J., I, Bernard, H., Ancrenaz, M., Coomes, D., Swinfield, T., Milodowski, D.T., Humle, T. & Struebig, M.J. (2019). Densities of Bornean orangutans (*Pongo pygmaeus morio*) in heavily degraded forest and oil palm plantations in Sabah, Borneo. *American Journal of Primatology*, 81(8): e23030. doi: 10.1002/ajp.23030.

137. Seavy, N.E., Viers, J.H. & Wood, J.K. (2009). Riparian bird response to vegetation structure: a multiscale analysis using LiDAR measurements of canopy height. *Ecological Applications*, 19(7): 1848-1857. doi: 10.1890/08-1124.1.
138. Seibold, S., Baessler, C., Brandl, R., Fahrig, L., Foerster, B., Heurich, M., Hothorn, T., Scheipl, F., Thorn, S. & Mueller, J. (2017). An experimental test of the habitat-amount hypothesis for saproxylic beetles in a forested region. *Ecology*, 98(6): 1613-1622. doi: 10.1002/ecy.1819.
139. Sillero, N. & Goncalves-Seco, L. (2014). Spatial structure analysis of a reptile community with airborne LiDAR data. *International Journal of Geographical Information Science*, 28(8): 1709-1722. doi: 10.1080/13658816.2014.902062.
140. Singh, M., Tokola, T., Hou, Z. & Notarnicola, C. (2017). Remote sensing-based landscape indicators for the evaluation of threatened-bird habitats in a tropical forest. *Ecology and Evolution*, 7(13): 4552-4567. doi: 10.1002/ece3.2970.
141. Singh, M., Cheyne, S.M. & Smith, D.A.E. (2018). How conspecific primates use their habitats: Surviving in an anthropogenically-disturbed forest in Central Kalimantan, Indonesia. *Ecological Indicators*, 87: 167-177. doi: 10.1016/j.ecolind.2017.12.041.
142. Smart, L.S., Swenson, J.J., Christensen, N.L. & Sexton, J.O. (2012). Three-dimensional characterization of pine forest type and red-cockaded woodpecker habitat by small-footprint, discrete-return lidar. *Forest Ecology and Management*, 281: 100-110. doi: 10.1016/j.foreco.2012.06.020.
143. Sovern, S.G., Lesmeister, D.B., Dugger, K.M., Pruett, M.S., Davis, R.J. & Jenkins, J.M. (2019). Activity center selection by northern spotted owls. *Journal of Wildlife Management*, 83(3): 714-727. doi: 10.1002/jwmg.21632.
144. Stephenson, N., Perroy, R., Eiben, J. & Klasner, F. (2017). High resolution habitat suitability modelling for an endemic restricted-range Hawaiian insect (*Nysius wekiuicola*, Hemiptera: Lygaeidae). *Journal of Insect Conservation*, 21(1): 87-96. doi: 10.1007/s10841-017-9956-4.
145. Svendsen, J.K., Sell, H., Bocher, P.K. & Svenning, J. (2015). Habitat and nest site preferences of Red-backed Shrike (*Lanius collurio*) in western Denmark. *Ornis Fennica*, 92(2): 63-75.
146. Swatantran, A., Dubayah, R., Goetz, S., Hofton, M., Betts, M.G., Sun, M., Simard, M. & Holmes, R. (2012). Mapping Migratory Bird Prevalence Using Remote Sensing Data Fusion. *Plos One*, 7(1): e28922. doi: 10.1371/journal.pone.0028922.
147. Swift, C.E., Vierling, K.T., Hudak, A.T. & Vierling, L.A. (2017). Relationships among Vegetation Structure, Canopy Composition, and Avian Richness Patterns across an Aspen-Conifer Forest Gradient. *Canadian Journal of Remote Sensing*, 43(3): 231-243. doi: 10.1080/07038992.2017.1317205.
148. Taillie, P.J., Moorman, C.E., Smart, L.S. & Pacifici, K. (2019). Bird community shifts associated with saltwater exposure in coastal forests at the leading edge of rising sea level. *Plos One*, 14(5): e0216540. doi: 10.1371/journal.pone.0216540.
149. Tattoni, C., Rizzoli, F. & Pedrini, P. (2012). Can LiDAR data improve bird habitat suitability models?. *Ecological Modelling*, 245: 103-110. doi: 10.1016/j.ecolmodel.2012.03.020.
150. Thers, H., Brunbjerg, A.K., Laessoe, T., Ejrnaes, R., Bocher, P.K. & Svenning, J. (2017). Lidar-derived variables as a proxy for fungal species richness and

- composition in temperate Northern Europe. *Remote Sensing of Environment*, 200: 102-113. doi: 10.1016/j.rse.2017.08.011.
151. Trainor, A.M., Walters, J.R., Morris, W.F., Sexton, J. & Moody, A. (2013). Empirical estimation of dispersal resistance surfaces: a case study with red-cockaded woodpeckers. *Landscape Ecology*, 28(4): 755-767. doi: 10.1007/s10980-013-9861-5.
  152. Tweedy, P.J., Moriarty, K.M., Bailey, J.D. & Epps, C.W. (2019). Using fine scale resolution vegetation data from LiDAR and ground-based sampling to predict Pacific marten resting habitat at multiple spatial scales. *Forest Ecology and Management*, 452: 117556. doi: 10.1016/j.foreco.2019.117556.
  153. Vauhkonen, J. & Imponen, J. (2016). Unsupervised classification of airborne laser scanning data to locate potential wildlife habitats for forest management planning. *Forestry*, 89(4): 350-363. doi: 10.1093/forestry/cpw011.
  154. Vierling, K.T., Baessler, C., Brandl, R., Vierling, L.A., Weiss, I. & Mueller, J. (2011). Spinning a laser web: predicting spider distributions using LiDAR. *Ecological Applications*, 21(2): 577-588. doi: 10.1890/09-2155.1.
  155. Vierling, K.T., Swift, C.E., Hudak, A.T., Vogeler, J.C. & Vierling, L.A. (2014). How much does the time lag between wildlife field- data collection and LiDAR- data acquisition matter for studies of animal distributions? A case study using bird communities. *Remote Sensing Letters*, 5(2): 185-193. doi: 10.1080/2150704X.2014.891773.
  156. Vierling, L.A., Vierling, K.T., Adam, P. & Hudak, A.T. (2013). Using Satellite and Airborne LiDAR to Model Woodpecker Habitat Occupancy at the Landscape Scale. *Plos One*, 8(12): e80988. doi: 10.1371/journal.pone.0080988.
  157. Vihervaara, P., Mononen, L., Auvinen, A., Virkkala, R., Lu, Y., Pippuri, I., Packalen, P., Valbuena, R. & Valkama, J. (2015). How to integrate remotely sensed data and biodiversity for ecosystem assessments at landscape scale. *Landscape Ecology*, 30(3): 501-516. doi: 10.1007/s10980-014-0137-5.
  158. Vogeler, J.C., Hudak, A.T., Vierling, L.A. & Vierling, K.T. (2013). Lidar-Derived Canopy Architecture Predicts Brown Creeper Occupancy of Two Western Coniferous Forests. *Condor*, 115(3): 614-622. doi: 10.1525/cond.2013.110082.
  159. Vogeler, J.C., Hudak, A.T., Vierling, L.A., Evans, J., Green, P. & Vierling, K.I.T. (2014). Terrain and vegetation structural influences on local avian species richness in two mixed-conifer forests. *Remote Sensing of Environment*, 147: 13-22. doi: 10.1016/j.rse.2014.02.006.
  160. Vogeler, J.C., Yang, Z. & Cohen, W.B. (2016). Mapping Suitable Lewis's Woodpecker Nesting Habitat in a Post-Fire Landscape. *Northwest Science*, 90(4): 421-432. doi: 10.3955/046.090.0404.
  161. Wallis, C.I.B., Paulsch, D., Zeilinger, J., Silva, B., Fernandez, G.F.C., Brandl, R., Farwig, N. & Bendix, J. (2016). Contrasting performance of Lidar and optical texture models in predicting avian diversity in a tropical mountain forest. *Remote Sensing of Environment*, 174: 223-232. doi: 10.1016/j.rse.2015.12.019.
  162. Weisberg, P.J., Dilts, T.E., Becker, M.E., Young, J.S., Wong-Kone, D.C., Newton, W.E. & Ammon, E.M. (2014). Guild-specific responses of avian species richness to LiDAR-derived habitat heterogeneity. *Acta Oecologica-International Journal of Ecology*, 59: 72-83. doi: 10.1016/j.actao.2014.06.002.

163. Work, T.T., St Onge, B. & Jacobs, J.M. (2011). Response of female beetles to LIDAR derived topographic variables in Eastern boreal mixedwood forests (Coleoptera, Carabidae). *Zookeys*(147): 623-639. doi: 10.3897/zookeys.147.2013.
164. Wu, Q., Lane, C. & Liu, H. (2014). An Effective Method for Detecting Potential Woodland Vernal Pools Using High-Resolution LiDAR Data and Aerial Imagery. *Remote Sensing*, 6(11): 11444-11467. doi: 10.3390/rs61111444.
165. Young, J.S., Ammon, E.M., Weisberg, P.J., Dilts, T.E., Newton, W.E., Wong-Kone, D.C. & Heki, L.G. (2013). Comparison of bird community indices for riparian restoration planning and monitoring. *Ecological Indicators*, 34: 159-167. doi: 10.1016/j.ecolind.2013.05.004.
166. Zabihi, K., Driese, K.L., Paige, G.B. & Hild, A.K. (2019). Application of Ground-Based Lidar and Gap Intercept Measurements to Quantify a Shrub Configuration Metric within Greater Sage-Grouse Nesting Habitat. *Western North American Naturalist*, 79(4): 500-514. doi: 10.3398/064.079.0404.
167. Zellweger, F., Braunisch, V., Baltensweiler, A. & Bollmann, K. (2013). Remotely sensed forest structural complexity predicts multi species occurrence at the landscape scale. *Forest Ecology and Management*, 307: 303-312. doi: 10.1016/j.foreco.2013.07.023.
168. Zellweger, F., Morsdorf, F., Purves, R.S., Braunisch, V. & Bollmann, K. (2014). Improved methods for measuring forest landscape structure: LiDAR complements field-based habitat assessment. *Biodiversity and Conservation*, 23(2): 289-307. doi: 10.1007/s10531-013-0600-7.
169. Zellweger, F., Braunisch, V., Morsdorf, F., Baltensweiler, A., Abegg, M., Roth, T., Bugmann, H. & Bollmann, K. (2015). Disentangling the effects of climate, topography, soil and vegetation on stand-scale species richness in temperate forests. *Forest Ecology and Management*, 349: 36-44. doi: 10.1016/j.foreco.2015.04.008.
170. Zellweger, F., Baltensweiler, A., Ginzler, C., Roth, T., Braunisch, V., Bugmann, H. & Bollmann, K. (2016). Environmental predictors of species richness in forest landscapes: abiotic factors versus vegetation structure. *Journal of Biogeography*, 43(6): 1080-1090. doi: 10.1111/jbi.12696.
171. Zellweger, F., Roth, T., Bugmann, H. & Bollmann, K. (2017). Beta diversity of plants, birds and butterflies is closely associated with climate and habitat structure. *Global Ecology and Biogeography*, 26(8): 898-906. doi: 10.1111/geb.12598.
172. Zhao, F., Sweitzer, R.A., Guo, Q. & Kelly, M. (2012). Characterizing habitats associated with fisher den structures in the Southern Sierra Nevada, California using discrete return lidar. *Forest Ecology and Management*, 280: 112-119. doi: 10.1016/j.foreco.2012.06.005.
173. Zielewska-Buettner, K., Heurich, M., Mueller, J. & Braunisch, V. (2018). Remotely Sensed Single Tree Data Enable the Determination of Habitat Thresholds for the Three-Toed Woodpecker (*Picoides tridactylus*). *Remote Sensing*, 10(12): 1972. doi: 10.3390/rs10121972.