

# Interindividual Differences in In Vitro Human Intestinal Microbial Conversion of 3-acetyl-DON and 15-acetyl-DON

Fangfang Li <sup>1,†</sup>, Jing Jin <sup>1,2,†</sup>, Ivonne M. C. M. Rietjens <sup>2</sup> and Fuguo Xing <sup>1,\*</sup>

<sup>1</sup> Institute of Food Science and Technology, Chinese Academy of Agricultural Sciences/Key Laboratory of Agro-Products Quality and Safety Control in Storage and Transport Process, Ministry of Agriculture and Rural Affairs of P. R. China, 2 Yuanmingyuan West Road, Haidian District, Beijing 100193, China; 1742240749@163.com (F.L.); jinjing@caas.cn (J.J.)


<sup>2</sup> Division of Toxicology, Wageningen University and Research, Stippeneng 4, 6708 WE Wageningen, The Netherlands; ivonne.rietjens@wur.nl

\* Correspondence: xingfuguo@caas.cn

† These authors contribute equally to this work.

**Table S1.** Information for the individual fecal samples collected in this study.

Individual	Gender	Age
1	M	26
2	M	25
3	F	25
4	F	25
5	F	25
6	F	24
7	F	25
8	F	26
9	F	25
10	M	30
11	M	26
12	F	24
13	F	26
14	F	25
15	M	27
16	F	25
17	F	25
18	F	25
19	F	25
20	M	26
21	F	25
22	F	25
23	F	25
24	F	25
25	F	24



**Figure S1.** Analysis of the microbiome relative abundance of species at the phylum level in individual human samples.