water

MDPI

Supplementary Material

Hydrological Response of the Kunhar River Basin in Pakistan to Climate Change and Anthropogenic Impacts on Runoff Characteristics

Muhammad Saifullah [1,*], Muhammad Adnan [2], Muhammad Zaman [3], Andrzej Wałęga [4], Shiyin Liu [2,*], Muhammad Imran Khan [5], Alexandre S. Gagnon [6] and Sher Muhammad [7]

1 Department of Agricultural Engineering, Muhammad Nawaz Shareef University of Agriculture, Multan 66000, Pakistan
2 Institute of International Rivers and Eco-Security, Yunnan University, Kunming 650500, China; adnan@lzb.ac.cn
3 Department of Irrigation and Drainage, University of Agriculture, Faisalabad 380000, Pakistan; muhammad.zaman@uaf.edu.pk
4 Department of Sanitary Engineering and Water Management, University of Agriculture in Krakow, 31-120 Kraków, Poland; andrzej.walega@urk.edu.pl
5 Research Center of Fluid Machinery Engineering and Technology, Jiangsu University, Zhenjiang 212013, China; imrankhan7792@yahoo.com
6 School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool L3 3AF, UK; A.Gagnon@ljmu.ac.uk
7 International Centre for Integrated Mountain Development (ICIMOD), Lalitpur 44700, Nepal; sher.muhammad@icimod.org
* Correspondence: muhammad.saifullah@mnsuam.edu.pk (M.S.); shiyin.liu@ynu.edu.cn (S.L.)



Figure S1. MK trend analysis of Runoff (R), Precipitation (Pr), and Evapotranspiration (Et) for whole period (1971–2010)



Figure S2. MK trend analysis of Precipitation (Pr), Evapotranspiration (Et), and Runoff (R) during pre-change (1971–1996) and post-change (1997–2010) periods



Figure S3. ITA of precipitation (**a**, **b**), evapotranspiration (**c**, **d**), and runoff (**e, f**) during pre-change (1971–1996), and post-change (1997–2010) periods. Note:* obseved data points